arXiv:1507.03834v2 [cs.SC] 8 Aug 2015

Open Weak CAD and Its Applications

Liyun Dai, Jingjun Han *

LMAM & School of Mathematical Sciences
& Beijing International Center for Mathematical Research
Peking University, Beijing 100871, China

Hoon Hong

Department of Mathematics, North Carolina State University, Raleigh NC 27695, USA

Bican Xia

LMAM & School of Mathematical Sciences, Peking University, Beijing 100871, China

---

Abstract

The concept of open weak CAD is introduced. Every open CAD is an open weak CAD. On the contrary, an open weak CAD is not necessarily an open CAD. An algorithm for computing open weak CADs is proposed. The key idea is to compute the intersection of projection factor sets produced by different projection orders. The resulting open weak CAD often has smaller number of sample points than open CADs.

The algorithm can be used for computing sample points for all open connected components of $f \neq 0$ for a given polynomial f. It can also be used for many other applications, such as testing semi-definiteness of polynomials and copositive problems. In fact, we solved several difficult semi-definiteness problems efficiently by using the algorithm. Furthermore, applying the algorithm to copositive problems, we find an explicit expression of the polynomials producing open weak CADs under some conditions, which significantly improves the efficiency of solving copositive problems.

Key words: Open weak CAD, open delineable, CAD projection, semi-definiteness, copositivity.

⋆ This research was partly supported by US National Science Foundation Grant 1319632 and National Science Foundation of China Grants 11290141 and 11271034.

* Corresponding author

Email addresses: `dailiyun@pku.edu.cn`, `hanjingjunfdfz@gmail.com` (Liyun Dai, Jingjun Han), `hong@ncsu.edu` (Hoon Hong), `xbc@math.pku.edu.cn` (Bican Xia).

URLs: `www.math.ncsu.edu/~hong` (Hoon Hong), `www.math.pku.edu.cn/is/~xbc/` (Bican Xia).

1. Introduction

The cylindrical algebraic decomposition (CAD) method was first proposed by Collins (1975, 1998). A key role in CAD algorithm is its projection operator. An improvement of CAD projection is Hong's projection operator (Hong, 1990). For many problems, a smaller projection operator given by McCallum (McCallum, 1988, 1998), with an improvement by Brown (Brown, 2001), is more efficient.

For a polynomial $f \in \mathbb{R}[x_1 \ldots, x_n]$, the graph of f decomposes $\mathbb{R}^n$ into finitely many connected regions with different dimensions. Typical CAD methods compute a CAD (represented by sample points) of $\mathbb{R}^n$ which has a cylindrical structure and guarantees that the intersection of the CAD (the set of sample points) and every region defined by f is not empty. It is well known that the scale (the number of sample points) of CAD is usually much bigger than the number of regions defined by the polynomial. It is also the case that the scale of *open CAD* (see Definition 14) is usually much larger than the number of open regions defined by $f \neq 0$.

In this paper, we introduce the concept of *open weak CAD*. Open weak CADs might not have cylindrical structure in the sense of classical CAD. However, open weak CADs preserve some geometric information, *i.e.*, they are *open weak delineable* (see Definition 1). As the result, every open CAD is an open weak CAD. On the contrary, an open weak CAD is not necessarily an open CAD.

An algorithm for computing open weak CADs is proposed. The key idea is to compute the intersection of projection factor sets produced by different projection orders. The resulting open weak CAD often has smaller number of sample points than open CADs.

The algorithm can be used for computing sample points for all open connected components of $f \neq 0$ for a given polynomial f. It can also be used for many other applications, such as testing semi-definiteness of polynomials and copositive problems. In fact, we solved several difficult semi-definiteness problems efficiently by using the algorithm. Furthermore, applying the algorithm to copositive problems, we find an explicit expression of the polynomials producing open weak CADs under some conditions, which significantly improves the efficiency of solving copositive problems.

The structure of this paper is as follows. In Section 2, the definitions of open weak delineable and open weak CAD are introduced and the problem studied in this paper is formally stated. In Section 3, we review basic definitions, lemmas and concepts of CAD. In Section 4, an algorithm for computing open weak CAD is proposed and is proved to be correct. In Section 5, as an application of the algorithm, we show how to compute open sample. In Section 6, as an application of the algorithm, a new method for polynomial inequality proving is proposed by combining our previous work (Han et al., 2013) with the algorithm in Section 4. Section 7 introduces another important application of the algorithm to testing copositive problem. Section 8 includes several examples which demonstrate the effectiveness of the algorithms.

Compared to our ISSAC'2014 paper, this paper is totally re-constructed by clarifying basic concepts and the problem from a new point of view. Therefore, Section 2 is totally re-written to start the paper by formally defining the problem studied in this paper. The subsequent sections are modified according to the new viewpoint. Some proofs are modified and added. All important new definitions and algorithms are followed by illustrative examples. Section 7 (copositive problem) is new with substantial contents and new results. Section 8 contains new examples.

2. Problem: Open weak CAD

In this section, we describe the problem studied in this paper. First, let us introduce the concept of *open weak delineable.*

Definition 1 (Open weak delineable)**.** Let S be an open set of $\mathbb{R}^j (1 \le j < n)$. The polynomial $f \in \mathbb{R}[x_1, \dots, x_n]$ is said to be *open weak delineable* on S if, for any maximal open connected set $U \subseteq \mathbb{R}^n$ defined by $f \neq 0$, we have

$$(S \times \mathbb{R}^{n-j}) \bigcap U \neq \emptyset \iff \forall \boldsymbol{\alpha} \in S,\ (\boldsymbol{\alpha} \times \mathbb{R}^{n-j}) \bigcap U \neq \emptyset.$$

Let $h \in \mathbb{R}[x_1, \dots, x_j]$ for $j < n$. We say the polynomial f is *open weak delineable* over h in $\mathbb{R}^j$, if f is open weak delineable on any open connected component of $h \neq 0$ in $\mathbb{R}^j$.

Note that if f is *analytic delineable* (Collins (1975)) on S then f is also *open weak delineable* on S. However the converse is not necessarily true.

Example 1. Let

$$f = x_1 - x_2^4 + 10x_2^3 - 35x_2^2 + 50x_2 - 24 \in \mathbb{R}[x_1, x_2].$$

The plot of $f = 0$ is given by



Fig. 1.

Note that f is analytically delineable and also open weak delineable on the set $(-\infty, -1)$. Note that f is *not* analytically delineable but *is* open weak delineable on the set $(-1, \infty)$. Note also that f is open weak delineable over $h = x_1 + 1$ in $\mathbb{R}$.

Definition 2 (Open weak CAD)**.** Let $f \in \mathbb{R}[x_1, \dots, x_n]$. A decomposition of $\mathbb{R}^j$, $1 \le j < n$, is called an *open weak CAD* of f in $\mathbb{R}^j$ if and only if f is open weak delineable on every j-dimensional open set in the decomposition.

Example 2. Let f be the polynomial from Example 1. Let $k = 1$. Then

$$\{(-\infty, -1), [-1, -1], (-1, \infty)\}$$

is an open weak CAD of f.

Finally, we are ready to state the problem precisely.

Problem. (Open Weak CAD) Devise an algorithm with the following specification.

In: $f \in \mathbb{Z}[x_1, \ldots, x_n]$

Out: $h_1, h_2, \ldots, h_{n-1}$ where $h_j \in \mathbb{Z}[x_1, \ldots, x_j]$ such that f is open weak delineable over h_j in $\mathbb{R}^j$.

We call the number of open components in $\mathbb{R}^j$ defined by $h_j \neq 0$, the *scale* of the open weak CAD of f defined by h_j in $\mathbb{R}^j$.

Remark 3. The output of the above problem is a list of "projection" polynomials, not an open weak CAD. However it is trivial to compute sample points in an open weak CAD of f from the projection polynomials. Thus, we will call the above problem "Open Weak CAD".

Example 3. Consider the following polynomial.

In: $f = (x_3^2 + x_2^2 + x_1^2 - 1)(4x_3 + 3x_2 + 2x_1 - 1) \in \mathbb{Z}[x_1, x_2, x_3]$

Out: $h_1 = (x_1-1)(x_1+1)(29x_1^2-4x_1-24)((20x_1^2-4x_1-15)^2+(13x_1^2-4x_1-8)^2) \in \mathbb{Z}[x_1]$

$h_2 = (x_2^2 + x_1^2 - 1)(25x_2^2 + 12x_2x_1 + 20x_1^2 - 6x_2 - 4x_1 - 15) \in \mathbb{Z}[x_1, x_2]$

The left plot in Figure 2 below shows the open weak CAD of f produced by h_1 and h_2. The factor $(20x_1^2 - 4x_1 - 15)^2 + (13x_1^2 - 4x_1 - 8)^2$ in h_1 does not have real root and thus it does not contribute to the open weak CAD.



Fig. 2. Example 3

Remark 4. For comparison, if we apply an open CAD algorithm on the above f, one would obtain the following output

Out: $h_1 = (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)(13x_1^2 - 4x_1 - 8) \in \mathbb{Z}[x_1]$

$h_2 = (x_2^2 + x_1^2 - 1)(25x_2^2 + 12x_2x_1 + 20x_1^2 - 6x_2 - 4x_1 - 15) \in \mathbb{Z}[x_1, x_2]$

The right plot in Figure 2 shows the open CAD of f produced by h_1 and h_2. Note that it has more cells than the open weak CAD (on the left).

Remark 5. It is natural to wonder whether the multivariate discriminants of f always produce open weak CADs. Unfortunately, this is not true since the discriminant $\mathrm{discrim}(f, [x_n, \ldots, x_{j+1}])$ may vanish identically and thus does not always produces an open weak CAD of $\mathbb{R}^j$. One may also wonder whether if the multivariate discriminants of f do produce open weak CADs, then they would be the smallest open weak CADs. Unfortunately this is not true, either. In Example 1, it has been shown that $x_1 + 1$ produces an open weak CAD of $\mathbb{R}$ with 2 open intervals. But the discriminant $\mathrm{discrim}(f, x_2) = -16(16x_1 - 9)(x_1 + 1)^2$ produces an open weak CAD of $\mathbb{R}$ with 3 open intervals.

3. Preliminaries

In this section, we review some basic (known) concepts and results. If not specified, for a positive integer n, let $\boldsymbol{x}_n$ be the list of variable $(x_1, \ldots, x_n)$ and $\boldsymbol{\alpha}_n$ and $\boldsymbol{\beta}_n$ denote the point $(\alpha_1, \ldots, \alpha_n) \in \mathbb{R}^n$ and $(\beta_1, \ldots, \beta_n) \in \mathbb{R}^n$, respectively.

Definition 6. Let $f \in \mathbb{Z}[\boldsymbol{x}_n]$, denote by $\mathrm{lc}(f, x_i)$ and $\mathrm{discrim}(f, x_i)$ the *leading coefficient* and the *discriminant* of f with respect to (w.r.t.) x_i, respectively. The set of real zeros of f is denoted by $\mathrm{Zero}(f)$. Denote by $\mathrm{Zero}(L)$ or $\mathrm{Zero}(f_1, \ldots, f_m)$ the common real zeros of $L = \{f_1, \ldots, f_m\} \subset \mathbb{Z}[\boldsymbol{x}_n]$. The *level* for f is the biggest j such that $\deg(f, x_j) > 0$ where $\deg(f, x_j)$ is the degree of f w.r.t. x_j. For polynomial set $L \subseteq \mathbb{Z}[\boldsymbol{x}_n]$, $L^{[i]}$ is the set of polynomials in L with level i.

Definition 7. Let P_n be the symmetric permutation group of $x_1, \ldots, x_n$. Define $P_{n,i}$ to be the subgroup of P_n, where any element σ of $P_{n,i}$ fixes $x_1, \ldots, x_{i-1}$, *i.e.*, $\sigma(x_j) = x_j$ for $j = 1, \ldots, i-1$.

Definition 8. If $h \in \mathbb{Z}[\boldsymbol{x}_n]$ can be factorized in $\mathbb{Z}[\boldsymbol{x}_n]$ as:

$$h = a l_1^{2j_1-1} \cdots l_t^{2j_t-1} {h_1}^{2i_1} \cdots {h_m}^{2i_m},$$

where $a \in \mathbb{Z}$, $t \geq 0, m \geq 0$, $l_j (i = 1, \ldots, t)$ and $h_i (i = 1, \ldots, m)$ are pairwise different irreducible primitive polynomials with positive leading coefficients (under a suitable ordering) and positive degrees in $\mathbb{Z}[\boldsymbol{x}_n]$, then define

$$\begin{aligned}
&\mathrm{sqrfree}(h) = l_1 \cdots l_t h_1 \cdots h_m,\\
&\mathrm{sqrfree}_1(h) = \{l_i, i = 1, 2, \ldots, t\},\\
&\mathrm{sqrfree}_2(h) = \{h_i, i = 1, 2, \ldots, m\}.
\end{aligned}$$

If h is a constant, let $\mathrm{sqrfree}(h) = 1$, $\mathrm{sqrfree}_1(h) = \mathrm{sqrfree}_2(h) = \{1\}$.

In the following, we introduce some basic concepts and results of CAD. The reader is referred to Collins (1975); Hong (1990); Collins and Hong (1991); McCallum (1988, 1998); Brown (2001) for a detailed discussion on the properties of CAD.

Definition 9. (Collins, 1975; McCallum, 1988) An n-variate polynomial $f(\boldsymbol{x}_{n-1}, x_n)$ over the reals is said to be *delineable* on a subset S (usually connected) of $\mathbb{R}^{n-1}$ if (1) the portion of the real variety of f that lies in the cylinder $S \times \mathbb{R}$ over S consists of the union of the graphs of some $t \geq 0$ continuous functions $\theta_1 < \cdots < \theta_t$ from S to $\mathbb{R}$; and (2) there exist integers $m_1, \ldots, m_t \geq 1$ s.t. for every $a \in S$, the multiplicity of the root $\theta_i(a)$ of $f(a, x_n)$ (considered as a polynomial in x_n alone) is m_i.

Definition 10. (Collins, 1975; McCallum, 1988) In the above definition, the θ_i are called the real root functions of f on S, the graphs of the θ_i are called the f-*sections* over S, and the regions between successive f-sections are called f-*sectors*.

Theorem 11. *(McCallum, 1988, 1998) Let $f(\boldsymbol{x}_n, x_{n+1})$ be a polynomial in $\mathbb{Z}[\boldsymbol{x}_n, x_{n+1}]$ of positive degree and* $\mathrm{discrim}(f, x_{n+1})$ *is a nonzero polynomial. Let S be a connected submanifold of $\mathbb{R}^n$ on which f is degree-invariant and does not vanish identically, and in which* $\mathrm{discrim}(f, x_{n+1})$ *is order-invariant. Then f is analytic delineable on S and is order-invariant in each f-section over S.*

Based on this theorem, McCallum proposed the projection operator MCproj, which consists of the discriminant of f and all coefficients of f.

Theorem 12. *(Brown, 2001) Let $f(\boldsymbol{x}_n, x_{n+1})$ be a $(n+1)$-variate polynomial of positive degree in x_{n+1} such that* $\mathrm{discrim}(f, x_{n+1}) \neq 0$. *Let S be a connected submanifold of $\mathbb{R}^n$ in which* $\mathrm{discrim}(f, x_{n+1})$ *is order-invariant, the leading coefficient of f is sign-invariant, and such that f vanishes identically at no point in S. f is degree-invariant on S.*

Based on this theorem, Brown obtained a reduced McCallum projection in which only leading coefficients, discriminants and resultants appear. The Brown projection operator is defined as follows.

Definition 13. (Brown, 2001) Given a polynomial $f \in \mathbb{Z}[\boldsymbol{x}_n]$, if f is with level n, the Brown projection operator for f is

$$\mathtt{Bp}(f, [x_n]) = \mathrm{Res}(\mathrm{sqrfree}(f), \frac{\partial(\mathrm{sqrfree}(f))}{\partial x_n}, x_n).$$

Otherwise $\mathtt{Bp}(f, [x_n]) = f$. If L is a polynomial set with level n, then

$$\begin{aligned}\mathtt{Bp}(L, [x_n]) = \bigcup_{f \in L} \{\mathrm{Res}(\mathrm{sqrfree}(f), \frac{\partial(\mathrm{sqrfree}(f))}{\partial x_n}, x_n)\} \bigcup \\ \bigcup_{f,g \in L, f \neq g} \{\mathrm{Res}(\mathrm{sqrfree}(f), \mathrm{sqrfree}(g), x_n)\}.\end{aligned}$$

Define

$$\begin{aligned}&\mathtt{Bp}(f, [x_n, x_{n-1}, \ldots, x_i]) \\ =&\mathtt{Bp}(\mathtt{Bp}(f, [x_n, x_{n-1}, \ldots, x_{i+1}]), [x_i]).\end{aligned}$$

The following definition of *open CAD* is essentially the GCAD introduced in Strzeboński (2000). For convenience, we use the terminology of open CAD in this paper.

Definition 14. (Open CAD) For a polynomial $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$, an open CAD defined by $f(\boldsymbol{x}_n)$ under the order $x_1 \prec x_2 \prec \cdots \prec x_n$ is a set of sample points in $\mathbb{R}^n$ obtained through the following three phases:

(1) [Projection] Use the Brown projection operator on $f(\boldsymbol{x}_n)$, let $F = \{f, \mathtt{Bp}(f, [x_n]), \ldots, \mathtt{Bp}(f, [x_n, \ldots, x_2])\}$;
(2) [Base] Choose one rational point in each of the open intervals defined by the real roots of $F^{[1]}$;
(3) [Lifting] Substitute each sample point of $\mathbb{R}^{i-1}$ for $\boldsymbol{x}_{i-1}$ in $F^{[i]}$ to get a univariate polynomial $F_i(x_i)$ and then, by the same method as Base phase, choose sample points for $F_i(x_i)$. Repeat the process for i from 2 to n.

Sometimes, we say that the polynomial set F produces the open CAD or simply, F is an open CAD.

4. Algorithm: Open weak CAD

In this section, we describe an algorithm (Algorithm 1) for computing open weak CAD and prove its correctness (Theorem 32). We begin by introducing a “projection” operator `Hp`.

Definition 15 (Open weak projection operator)**.** Let $f \in \mathbb{Z}[x_1, \ldots, x_n]$. For given $m (1 \le m \le n)$, denote $[\boldsymbol{y}] = [y_1, \ldots, y_m]$ where $y_i \in \{x_1, \ldots, x_n\}$ for $1 \le i \le m$ and $y_i \ne y_j$ for $i \ne j$. For $1 \le i \le m$, $\mathtt{Hp}(f, [\boldsymbol{y}], y_i)$ and $\mathtt{Hp}(f, [\boldsymbol{y}])$ are defined recursively as follows.

$$\begin{aligned} \mathtt{Bp}(f, [y_i]) &= \mathrm{Res}(\mathrm{sqrfree}(f), \frac{\partial(\mathrm{sqrfree}(f))}{\partial y_i}, y_i) \\ \mathtt{Hp}(f, [\boldsymbol{y}], y_i) &= \mathtt{Bp}(\mathtt{Hp}(f, \hat{[\boldsymbol{y}]}_i), [y_i]), \\ \mathtt{Hp}(f, [\boldsymbol{y}]) &= \mathrm{gcd}(\mathtt{Hp}(f, [\boldsymbol{y}], y_1), \ldots, \mathtt{Hp}(f, [\boldsymbol{y}], y_m)), \\ \mathtt{Hp}(f, [\,]) &= f, \end{aligned}$$

where $\hat{[\boldsymbol{y}]}_i = [y_1, \ldots, y_{i-1}, y_{i+1}, \ldots, y_m]$.

Example 4. We have

$$\begin{aligned} \mathtt{Hp}(f, [x_1, x_2]) &= \mathrm{gcd}\left(\mathtt{Hp}\left(f, [x_1, x_2], x_1\right), \mathtt{Hp}\left(f, [x_1, x_2], x_2\right)\right) \\ \mathtt{Hp}\left(f, [x_1, x_2], x_1\right) &= \mathtt{Bp}(\mathtt{Hp}(f, [x_2]), [x_1]) \\ \mathtt{Hp}\left(f, [x_1, x_2], x_2\right) &= \mathtt{Bp}(\mathtt{Hp}(f, [x_1]), [x_2]) \\ \mathtt{Hp}(f, [x_2]) &= \mathtt{Hp}(f, [x_2], x_2) \\ \mathtt{Hp}(f, [x_1]) &= \mathtt{Hp}(f, [x_1], x_1) \\ \mathtt{Hp}(f, [x_2], x_2) &= \mathrm{gcd}(\mathtt{Bp}(\mathtt{Hp}(f, [\,]), [x_2])) \\ \mathtt{Hp}(f, [x_1], x_1) &= \mathrm{gcd}(\mathtt{Bp}(\mathtt{Hp}(f, [\,]), [x_1])) \\ \mathtt{Hp}(f, [\,]) &= f \end{aligned}$$

Condensing the above expressions, we have

$$\mathtt{Hp}(f, [x_1, x_2]) = \mathrm{gcd}\left(\mathtt{Bp}(\mathtt{Bp}(f, [x_2]), [x_1]), \mathtt{Bp}(\mathtt{Bp}(f, [x_1]), [x_2])\right)$$

Algorithm 1. `Open Weak CAD`

```
Input: $f \in \mathbb{Z}[x_1, \dots, x_n]$
Output: $h_1, h_2, \dots, h_{n-1}$ where $h_j \in \mathbb{Z}[x_1, \dots, x_j]$ such that each $h_j$ produces an open weak CAD of $\mathbb{R}^j$ w.r.t. $f$
1: For all $1 \le j < n$ and $j < t \le n$, compute $\texttt{Hp}(f, [x_n, \dots, x_{j+1}], x_t)$ by Definition 15.
2: Compute $h_1(f), h_2(f), \dots, h_{n-1}(f)$ where
       $h_j(f) = \text{sqrfree}\left(\sum_{t=j+1}^{n} \texttt{Hp}(f, [x_n, \dots, x_{j+1}], x_t)^2\right).$
```

The Algorithm 1 (`Open Weak CAD`) based on the new operator `Hp` solves the problem proposed in Section 2.

Example 5. We illustrate the Algorithm 1 using the polynomial f from Example 3.

In: $f = (x_3^2 + x_2^2 + x_1^2 - 1)(4x_3 + 3x_2 + 2x_1 - 1) \in \mathbb{Z}[x_1, x_2, x_3]$

1: For all $1 \le j < 3$ and $j < t \le 3$, compute $\texttt{Hp}(f, [x_3, \dots, x_{j+1}], x_t)$ by Definition 15.

$$\begin{aligned}
\texttt{Hp}(f, [x_3], x_3) &= \texttt{Bp}(\texttt{Hp}(f, [\ ]), [x_3]) = \texttt{Bp}(f, [x_3]) \\
&= (x_2^2 + x_1^2 - 1)(25x_2^2 + 12x_2x_1 + 20x_1^2 - 6x_2 - 4x_1 - 15) \\
\texttt{Hp}(f, [x_3, x_2], x_2) &= \texttt{Bp}(\texttt{Hp}(f, [x_3]), [x_2]) = \texttt{Bp}(\texttt{Hp}(f, [x_3], x_3), [x_2]) \\
&= (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)(13x_1^2 - 4x_1 - 8) \\
\texttt{Hp}(f, [x_3, x_2], x_3) &= \texttt{Bp}(\texttt{Hp}(f, [x_2]), [x_3]) = \texttt{Bp}(\texttt{Hp}(f, [x_2], x_2), [x_3]) \\
&= (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)(20x_1^2 - 4x_1 - 15).
\end{aligned}$$

2: Compute $h_1(f), h_2(f)$ where

$$h_j(f) = \text{sqrfree}\left(\sum_{t=j+1}^{n} \texttt{Hp}(f, [x_3, \dots, x_{j+1}], x_t)^2\right).$$

$$\begin{aligned}
h_1 &= \text{sqrfree}((\texttt{Hp}(f, [x_3, x_2], x_2))^2 + (\texttt{Hp}(f, [x_3, x_2], x_3))^2) \\
&= (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)((20x_1^2 - 4x_1 - 15)^2 + (13x_1^2 - 4x_1 - 8)^2) \\
h_2 &= \text{sqrfree}((\texttt{Hp}(f, [x_3], x_3))^2) \\
&= (x_2^2 + x_1^2 - 1)(25x_2^2 + 12x_2x_1 + 20x_1^2 - 6x_2 - 4x_1 - 15)
\end{aligned}$$

Out: $h_1 = (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)((20x_1^2 - 4x_1 - 15)^2 + (13x_1^2 - 4x_1 - 8)^2)$

$h_2 = (x_2^2 + x_1^2 - 1)(25x_2^2 + 12x_2x_1 + 20x_1^2 - 6x_2 - 4x_1 - 15)$

Remark 16. Although H_j is more complicated than any $\texttt{Hp}(f, [x_n, \dots, x_{j+1}], x_t)$,

$$\text{Zero}(H_j) \subseteq \text{Zero}(\texttt{Hp}(f, [x_n, \dots, x_{j+1}], x_{j+1})) \subseteq \text{Zero}(\texttt{Bp}(f, [x_n, \dots, x_{j+1}])).$$

That means, for every open cell C' of open CAD produced by Brown's operator, there exists an open cell C of open weak CAD produced by H_j such that $C' \subseteq C$. Thus, the scale of open weak CAD is no bigger than that of open CAD.

Remark 17. In Algorithm 1, the scale of the open weak CAD of f defined by h_j in $\mathbb{R}^j$ is not always the smallest. For example, let f be the polynomial in Example 1, then $h_1 = (16x_1 - 9)(x_1 + 1)$, and f is open weak delineable over $x_1 + 1$, as mentioned earlier.

We now prove the correctness of Algorithm 1.

Definition 18. (Open sample) A set of sample points $S_f \subseteq \mathbb{R}^k \setminus \mathrm{Zero}(f)$ is said to be an *open sample* defined by $f(\boldsymbol{x}_k) \in \mathbb{Z}[\boldsymbol{x}_k]$ in $\mathbb{R}^k$ if it has the following property: for every open connected set $U \subseteq \mathbb{R}^k$ defined by $f \neq 0$, $S_f \cap U \neq \emptyset$.

Suppose $g(\boldsymbol{x}_k)$ is another polynomial. If S_f is an open sample defined by $f(\boldsymbol{x}_k)$ in $\mathbb{R}^k$ such that $g(\boldsymbol{\alpha}) \neq 0$ for any $\boldsymbol{\alpha} \in S_f$, then we denote the open sample by $S_{f,g}$.

As a corollary of Theorems 11 and 12, a property of open CAD (or GCAD) is that at least one sample point can be taken from every highest dimensional cell via the open CAD (or GCAD) lifting phase. So, an open CAD is indeed an open sample.

Obviously, there are various efficient ways to compute $S_{f,g}$ for two given *univariate* polynomials $f, g \in \mathbb{Z}[x]$. For example, we may choose one rational point from every open interval defined by the real zeros of f such that g does not vanish at this point. Therefore, we only describe the specification of such algorithms `SPOne` here and omit the details of the algorithms.

Definition 19. Let $\boldsymbol{\alpha}_j = (\alpha_1, \ldots, \alpha_j) \in \mathbb{R}^j$ and $S \subseteq \mathbb{R}$ be a finite set, define

$$\boldsymbol{\alpha}_j \boxplus S = \{(\alpha_1, \ldots, \alpha_j, \beta) \mid \beta \in S\}.$$

Algorithm 2. `SPOne`

```
Input: Two univariate polynomials $f, g \in \mathbb{Z}[x]$
Output: $S_{f,g}$, an open sample defined by $f(x)$ in $\mathbb{R}$ such that $g(\boldsymbol{\alpha}) \neq 0$ for any $\boldsymbol{\alpha} \in S_{f,g}$
```

Algorithm 3. `OpenSP`

```
Input: Two lists of polynomials $L_1 = [f_n(\boldsymbol{x}_n), \ldots, f_j(\boldsymbol{x}_j)]$, $L_2 = [g_n(\boldsymbol{x}_n), \ldots, g_j(\boldsymbol{x}_j)]$, and
    a set of points $S$ in $\mathbb{R}^j$
Output: A set of sample points in $\mathbb{R}^n$
$O := S$
for $i$ from $j + 1$ to $n$ do
    $P := \emptyset$
    for $\boldsymbol{\alpha}$ in $O$ do
        $P := P \bigcup (\boldsymbol{\alpha} \boxplus \mathtt{SPOne}(f_i(\boldsymbol{\alpha}, x_i), g_i(\boldsymbol{\alpha}, x_i)))$
    end for
    $O := P$
end for
return $O$
```

Remark 20. For a polynomial $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$, let

$$\begin{aligned} B_1 &= [f, \mathtt{Bp}(f, [x_n]), \ldots, \mathtt{Bp}(f, [x_n, \ldots, x_2])], \\ B_2 &= [1, \ldots, 1], \\ S &= \mathtt{SPOne}(\mathtt{Bp}(f, [x_n, \ldots, x_2]), 1), \end{aligned}$$

then $\mathtt{OpenSP}(B_1, B_2, S)$ is an open CAD (an open sample) defined by $f(\boldsymbol{x}_n)$.

We will provide in this section a method which computes two lists C_1 and C_2 where the polynomials in C_1 are factors of corresponding polynomials in B_1 and will prove that any $\mathtt{OpenSP}(C_1, C_2, S_{f_j,g_j})$ is an open sample of $\mathbb{R}^n$ defined by $f(\boldsymbol{x}_n)$ for any open sample S_{f_j,g_j} in $\mathbb{R}^j$ where $f_j \in C_1$ and $g_j \in C_2$.

Remark 21. The output of $\mathtt{OpenSP}(L_1, L_2, S)$ is dependent on the method of choosing sample points in Algorithm `SPOne`. In the following, when we use the terminology "any $\mathtt{OpenSP}(L_1, L_2, S)$", we mean "no matter which method is used in Algorithm `SPOne` for choosing sample points".

Definition 22. (Open delineable) Let

$$L_1 = [f_n(\boldsymbol{x}_n), f_{n-1}(\boldsymbol{x}_{n-1}), \ldots, f_j(\boldsymbol{x}_j)], \tag{1}$$

$$L_2 = [g_n(\boldsymbol{x}_n), g_{n-1}(\boldsymbol{x}_{n-1}), \ldots, g_j(\boldsymbol{x}_j)] \tag{2}$$

be two polynomial lists, S an open set of $\mathbb{R}^s$ $(s \le j)$ and $S' = S \times \mathbb{R}^{j-s}$. The polynomial $f_n(\boldsymbol{x}_n)$ is said to be *open delineable* on S w.r.t. L_1 and L_2, if $\mathcal{A} \bigcap U \neq \emptyset$ for any maximal open connected set $U \subseteq \mathbb{R}^n$ defined by $f_n \neq 0$ with $U \bigcap (S' \times \mathbb{R}^{n-j}) \neq \emptyset$ and any $\mathcal{A} = \mathtt{OpenSP}(L_1, L_2, \{\boldsymbol{\alpha}\})$ where $\boldsymbol{\alpha} \in S'$ is any point such that $f_j(\boldsymbol{\alpha}) g_j(\boldsymbol{\alpha}) \neq 0$.

Remark 23. Let $s = j$ in Definition 22, it could be shown that if $f_n(\boldsymbol{x}_n)$ is open delineable on S' w.r.t. L_1 and L_2, then $f_n(\boldsymbol{x}_n)$ is open weak delineable on $S' \backslash \mathrm{Zero}\{f_j g_j\}$.

Suppose $f_n(\boldsymbol{x}_n)$ is a squarefree polynomial in $\mathbb{Z}[\boldsymbol{x}_n]$ of positive degree and $S \subseteq \mathbb{R}^{n-1}$ is an open connected set in which $\mathtt{Bp}(f_n, [x_n])$ is sign-invariant. According to Theorem 11 and Theorem 12, f_n is analytic delineable on S. It is easy to see that f_n is open delineable on S w.r.t. $[f_n, \mathtt{Bp}(f_n, [x_n])]$ and $[f_n, \mathtt{Bp}(f_n, [x_n])]$.

Open delineability has the following four properties.

Proposition 24. (*open sample property*) *Let L_1, L_2 be as in Definition 22. If $f_n(\boldsymbol{x}_n)$ is open delineable on every open connected set of $f_j(\boldsymbol{x}_j) \neq 0$ w.r.t. L_1 and L_2, then for any open sample S_{f_j,g_j} in $\mathbb{R}^j$, any $\mathcal{A} = \mathtt{OpenSP}(L_1, L_2, S_{f_j,g_j})$ is an open sample defined by $f_n(\boldsymbol{x}_n)$ in $\mathbb{R}^n$.*

Proof. For any open connected set $U \subseteq \mathbb{R}^n$ defined by $f_n \neq 0$, there exists at least one open connected set $S \subseteq \mathbb{R}^j$ defined by $f_j \neq 0$ such that $U \bigcap (S \times \mathbb{R}^{n-j}) \neq \emptyset$. Since f_n is open delineable on S w.r.t. L_1 and L_2, we have $\mathcal{A} \bigcap U \neq \emptyset$ for any $\mathcal{A} = \mathtt{OpenSP}(L_1, L_2, S_{f_j,g_j})$. $\square$

Proposition 25. (*transitive property*) *Let L_1, L_2, S, S' be as in Definition 22. Suppose that there exists $k (j \le k \le n)$ such that $f_k(\boldsymbol{x}_k)$ is open delineable on S w.r.t. $L_1'' = [f_k(\boldsymbol{x}_k), \ldots, f_j(\boldsymbol{x}_j)]$ and $L_2'' = [g_k(\boldsymbol{x}_k), \ldots, g_j(\boldsymbol{x}_j)]$, and $f_n(\boldsymbol{x}_n)$ is open delineable on every open connected set of $f_k(\boldsymbol{x}_k) \neq 0$ w.r.t. $L_1' = [f_n(\boldsymbol{x}_n), \ldots, f_k(\boldsymbol{x}_k)]$ and $L_2' = [g_n(\boldsymbol{x}_n), \ldots, g_k(\boldsymbol{x}_k)]$. Then $f_n(\boldsymbol{x}_n)$ is open delineable on S w.r.t. L_1 and L_2.*

Proof. Let $\boldsymbol{\alpha} \in S'$ be any point such that $f_j(\boldsymbol{\alpha})g_j(\boldsymbol{\alpha}) \neq 0$, for any $\mathcal{A} = \texttt{OpenSP}(L_1, L_2, \{\boldsymbol{\alpha}\})$, we have $\mathcal{A} = \texttt{OpenSP}(L_1', L_2', \mathcal{A}')$ where $\mathcal{A}' = \texttt{OpenSP}(L_1'', L_2'', \{\boldsymbol{\alpha}\})$. For any open connected set $U \subseteq \mathbb{R}^n$ defined by $f_n \neq 0$ with $U \bigcap (S' \times \mathbb{R}^{n-j}) \neq \emptyset$, there exists an open connected set $V \subseteq \mathbb{R}^k$ defined by $f_k \neq 0$ with $U \bigcap (V \times \mathbb{R}^{n-k}) \neq \emptyset$ and $V \bigcap (S \times \mathbb{R}^{k-s}) \neq \emptyset$. Now we have $\mathcal{A}' \bigcap V \neq \emptyset$ since $f_k(\boldsymbol{x}_k)$ is open delineable on S w.r.t. L_1'' and L_2''. And then, $\mathcal{A} \bigcap U \neq \emptyset$ is implied by $U \bigcap (V \times \mathbb{R}^{n-k}) \neq \emptyset$ since $f_n(\boldsymbol{x}_n)$ is open delineable on V w.r.t. L_1' and L_2'. □

Proposition 26. (*nonempty intersection property*) *Let L_1, L_2 be as in Definition 22. For two open sets S_1 and S_2 of $\mathbb{R}^s$ $(s \leq j)$ with $S_1 \bigcap S_2 \neq \emptyset$, if $f_n(\boldsymbol{x}_n)$ is open delineable on both S_1 and S_2 w.r.t. L_1 and L_2, then $f_n(\boldsymbol{x}_n)$ is open delineable on $S_1 \bigcup S_2$ w.r.t. L_1 and L_2.*

Proof. For any $\boldsymbol{\alpha}_1 \in S_1$, $\boldsymbol{\alpha}_2 \in S_2$, $\boldsymbol{\alpha}_3 \in S_1 \bigcap S_2$ with $f_j(\boldsymbol{\alpha}_i)g_j(\boldsymbol{\alpha}_i) \neq 0$, any $\mathcal{A}_i = \texttt{OpenSP}(L_1, L_2, \{\boldsymbol{\alpha}_i\})$, and open connected set $U \subseteq \mathbb{R}^n$ defined by $f_n \neq 0$, we have $U \bigcap (S_1 \times \mathbb{R}^{n-s}) \neq \emptyset \iff \mathcal{A}_1 \bigcap U \neq \emptyset \iff \mathcal{A}_3 \bigcap U \neq \emptyset \iff \mathcal{A}_2 \bigcap U \neq \emptyset \iff U \bigcap (S_2 \times \mathbb{R}^{n-s}) \neq \emptyset$. □

Proposition 27. (*union property*) *Let L_1, L_2 be as in Definition 22. For $\sigma \in P_{n,j+1}$, denote $\boldsymbol{y}_n = (y_1, \ldots, y_n) = \sigma(\boldsymbol{x}_n)$ and $\boldsymbol{y}_i = (y_1, \ldots, y_i)$. Let $L_1' = [f_n(\boldsymbol{x}_n),\ p_{n-1}(\boldsymbol{y}_{n-1}),\ \ldots,\ p_j(\boldsymbol{y}_j)]$ and $L_2' = [q_n(\boldsymbol{x}_n),\ q_{n-1}(\boldsymbol{y}_{n-1}),\ \ldots,\ q_j(\boldsymbol{y}_j)]$ where $p_i(\boldsymbol{y}_i)$ and $q_i(\boldsymbol{y}_i)$ are polynomials in i variables.*

For two open sets S_1 and S_2 of $\mathbb{R}^j$, if (a) $f_n(\boldsymbol{x}_n)$ is open delineable on both S_1 and S_2 w.r.t. L_1 and L_2, (b) $f_n(\boldsymbol{x}_n)$ is open delineable on $S_1 \bigcup S_2$ w.r.t. L_1' and L_2', and (c) $p_j(\boldsymbol{y}_j)q_j(\boldsymbol{y}_j)$ vanishes at no points in $S_1 \bigcup S_2$, then $f_n(\boldsymbol{x}_n)$ is open delineable on $S_1 \bigcup S_2$ w.r.t. L_1 and L_2.

Proof. Let $\boldsymbol{\alpha}_1 \in S_1, \boldsymbol{\alpha}_2 \in S_2$ be two points such that $g_j p_j q_j(\boldsymbol{\alpha}_t) \neq 0$ for $t = 1, 2$. Let $\mathcal{A}_t = \texttt{OpenSP}(L_1, L_2, \{\boldsymbol{\alpha}_t\})$ and $\mathcal{A}'_t = \texttt{OpenSP}(L_1', L_2', \{\boldsymbol{\alpha}_t\})$. For any open connected set U defined by $f_n \neq 0$ with $U \bigcap (\boldsymbol{\alpha}_1 \times \mathbb{R}^{n-j}) \neq \emptyset$, then $\mathcal{A}_1 \bigcap U \neq \emptyset$ and $\mathcal{A}'_1 \bigcap U \neq \emptyset$. Since $f_n(\boldsymbol{x}_n)$ is open delineable on $S_1 \bigcup S_2$ w.r.t. L_1' and L_2', we have $\mathcal{A}'_2 \bigcap U \neq \emptyset$ which implies that $U \bigcap (S_2 \times \mathbb{R}^{n-j}) \neq \emptyset$ and $\mathcal{A}_2 \bigcap U \neq \emptyset$. Therefore, $f_n(\boldsymbol{x}_n)$ is open delineable on $(S_1 \bigcup S_2) \backslash \mathrm{Zero}(p_j q_j)$ w.r.t. L_1 and L_2. Since $p_j q_j$ does not vanish at any point of $S_1 \bigcup S_2$, $f_n(\boldsymbol{x}_n)$ is open delineable on $S_1 \bigcup S_2$ w.r.t. L_1 and L_2. □

Example 6. We illustrate Proposition 27 using the polynomial f from Example 3.

$$f = (x_3^2 + x_2^2 + x_1^2 - 1)(4x_3 + 3x_2 + 2x_1 - 1) \in \mathbb{Z}[x_1, x_2, x_3]$$

Let

$$\begin{aligned}
L_1 &= [f, f_{x_3}, f_{x_3x_2}] \\
L_1' &= [f, f_{x_2}, f_{x_2x_3}] \\
L_2 &= L_1 \\
L_2' &= L_1'
\end{aligned}$$

where

$$\begin{aligned}
f_{x_3} &= (x_2^2 + x_1^2 - 1)(25x_2^2 + 12x_2x_1 + 20x_1^2 - 6x_2 - 4x_1 - 15),\\
f_{x_3x_2} &= (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)(13x_1^2 - 4x_1 - 8),\\
f_{x_2} &= (x_3^2 + x_1^2 - 1)(25x_3^2 + 16x_3x_1 + 13x_1^2 - 8x_3 - 4x_1 - 8),\\
f_{x_2x_3} &= (x_1 - 1)(x_1 + 1)(29x_1^2 - 4x_1 - 24)(20x_1^2 - 4x_1 - 15).
\end{aligned}$$

Let

$$S_1 = \left(\frac{2}{13} - \frac{6\sqrt{3}}{13}, \frac{2}{13} + \frac{6\sqrt{3}}{13}\right)$$

$$S_2 = \left(\frac{2}{13} + \frac{6\sqrt{3}}{13}, \frac{1}{10} + \frac{\sqrt{19}}{5}\right)$$

be two open intervals in x_1-axis, where $x_1 = \frac{2}{13} + \frac{6\sqrt{3}}{13}$ is one of the real roots of the equation $f_{x_3x_2} = 0$. By typical CAD methods, S_1 and S_2 are two different cells in x_1-axis.

It could be deduced easily by Theorem 11, Theorem 12, and Proposition 25 that the conditions (a) and (b) of Proposition 27 are satisfied. Since $f_{x_2x_3}$ vanishes at no points in $S_1 \bigcup S_2$, condition (c) is also satisfied.

By Proposition 27, f is open delineable on $S_1 \bigcup S_2$ w.r.t. L_1 and L_2. Roughly speaking, the real root of $x_1 - \frac{2}{13} - \frac{6\sqrt{3}}{13}$ would not affect the open delineability, thus we could combine the two cells S_1 and S_2.

For the same reason, the real roots of

$$(13x_1^2 - 4x_1 - 8)(20x_1^2 - 4x_1 - 15)$$

would not affect the open delineability either.

Lemma 28. *(Han et al., 2013) Let f and g be coprime in $\mathbb{Z}[\boldsymbol{x}_n]$. For any connected open set U of $\mathbb{R}^n$, the open set $V = U \backslash \mathrm{Zero}(f, g)$ is also connected.*

Lemma 29. *Let $f = \gcd(f_1, \ldots, f_m)$ where $f_i \in \mathbb{Z}[\boldsymbol{x}_n]$, $i = 1, 2, \ldots, m$. Suppose f has no real zeros in a connected open set $U \subseteq \mathbb{R}^n$, then the open set $V = U \backslash \mathrm{Zero}(f_1, \ldots, f_m)$ is also connected.*

Proof. Without loss of generality, we can assume that $f = 1$. If $m = 1$, the result is obvious. The result of case $m = 2$ is just the claim of Lemma 28. For $m \geq 3$, let $g = \gcd(f_1, \ldots, f_{m-1})$ and $g_i = f_i/g$ $(i = 1, \ldots, m-1)$, then $\gcd(f_m, g) = 1$ and $\gcd(g_1, \ldots, g_{m-1}) = 1$. Let $A = \mathrm{Zero}(f_1, \ldots, f_m)$, $B = \mathrm{Zero}(g_1, \ldots, g_{m-1}) \bigcup \mathrm{Zero}(g, f_m)$. Since $A \subseteq B$, we have $U \backslash B \subseteq U \backslash A$. Notice that the closure of $U \backslash B$ equals the closure of $U \backslash A$, it suffices to prove that $U \backslash B$ is connected, which follows directly from Lemma 28 and induction. □

Definition 30. Define

$$\begin{aligned}
\overline{\mathtt{Hp}}(f, i) &= \{f, \mathtt{Hp}(f, [x_n]), \ldots, \mathtt{Hp}(f, [x_n, \ldots, x_i])\},\\
\widetilde{\mathtt{Hp}}(f, i) &= \{f, \mathtt{Hp}(f, [x_n], x_n), \ldots, \mathtt{Hp}(f, [x_n, \ldots, x_i], x_i)\}.
\end{aligned}$$

As a Corollary of Theorem 11 and Theorem 12, we have

Proposition 31. *Let* $f \in \mathbb{Z}[\boldsymbol{x}_n]$ *be a squarefree polynomial with level* n. *Then* f *is open delineable on every open connected set defined by* $\mathtt{Bp}(f,[x_n]) \neq 0$ *in* $\mathbb{R}^{n-1}$ *w.r.t.* $\overline{\mathtt{Hp}}(f,n)$ *and* $\widetilde{\mathtt{Hp}}(f,n)$.

The following Theorem is the main result of this paper, which guarantees the correctness of Algorithm 1.

Theorem 32. *Let* j *be an integer and* $2 \leq j \leq n$. *For any given polynomial* $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$ *and any open connected set* $U \subseteq \mathbb{R}^{j-1}$ *of* $\mathtt{Hp}(f,[x_n,\ldots,x_j]) \neq 0$, *let* $S = U \backslash \mathrm{Zero}(\{\mathtt{Hp}(f,[x_n,\ldots,x_j],x_t) \mid t = j,\ldots,n\})$. *Then* $f(\boldsymbol{x}_n)$ *is open delineable on the open connected set* S *w.r.t.* $\overline{\mathtt{Hp}}(f,j)$ *and* $\widetilde{\mathtt{Hp}}(f,j)$. *As a result,* f *is open weak delineable over* $H_{j-1}(f)$ *in* $\mathbb{R}^{j-1}$, *and Algorithm 1 is correct.*

Proof. First, by Lemma 29, S is open connected. We prove the theorem by induction on $k = n - j$. When $k = 0$, it is obviously true from Proposition 31. Suppose the theorem is true for all polynomials $g(\boldsymbol{x}_k) \in \mathbb{Z}[\boldsymbol{x}_k]$ with $k = 0,1,\ldots,n-i-1$. We now consider the case $k = n - i$. Let $[\boldsymbol{z}] = [x_n,\ldots,x_i]$. For any given polynomial $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$, let $U \subseteq \mathbb{R}^{i-1}$ be an open connected set of $\mathtt{Hp}(f,[\boldsymbol{z}]) \neq 0$ and $S = U \backslash \mathrm{Zero}(\{\mathtt{Hp}(f,[\boldsymbol{z}],x_t) \mid t = i,\ldots,n\})$.

For any point $\boldsymbol{\alpha} \in S$ with $\mathtt{Hp}(f,[\boldsymbol{z}],x_i)(\boldsymbol{\alpha}) \neq 0$, there exists an open connected set $S_{\boldsymbol{\alpha}} \subseteq \mathbb{R}^{i-1}$ such that $\boldsymbol{\alpha} \in S_{\boldsymbol{\alpha}}$ and $0 \notin \mathtt{Hp}(f,[\boldsymbol{z}],x_i)(S_{\boldsymbol{\alpha}})$. By induction, $\mathtt{Hp}(f,[x_n,\ldots,x_{i+1}])$ is open delineable on $S_{\boldsymbol{\alpha}}$ w.r.t. $\{\mathtt{Hp}(f,[\boldsymbol{z}])\}$ and $\{\mathtt{Hp}(f,[\boldsymbol{z}],x_i)\}$. By induction again and the transitive property of open delineable (Proposition 25), f is open delineable on $S_{\boldsymbol{\alpha}}$ w.r.t. $\overline{\mathtt{Hp}}(f,i)$ and $\widetilde{\mathtt{Hp}}(f,i)$.

For any point $\boldsymbol{\alpha} \in S$ with $\mathtt{Hp}(f,[\boldsymbol{z}],x_i)(\boldsymbol{\alpha}) = 0$, there exists an i' such that $n \geq i' \geq i+1$ and $\mathtt{Hp}(f,[\boldsymbol{z}],x_{i'})(\boldsymbol{\alpha}) \neq 0$. Thus there exists an open connected set $S'_{\boldsymbol{\alpha}}$ of $\mathbb{R}^{i-1}$ such that $\boldsymbol{\alpha} \in S'_{\boldsymbol{\alpha}}$ and $0 \notin \mathtt{Hp}(f,[\boldsymbol{z}],x_{i'})(S'_{\boldsymbol{\alpha}})$. Let $\sigma \in P_{n,i}$ with $\sigma(x_i) = x_{i'}$, in such case, $f(\sigma(\boldsymbol{x}_n))$ is open delineable on $S'_{\boldsymbol{\alpha}}$ w.r.t. $\overline{\mathtt{Hp}}(f(\sigma(\boldsymbol{x}_n)),i)$ and $\widetilde{\mathtt{Hp}}(f(\sigma(\boldsymbol{x}_n)),i)$. For any $\boldsymbol{\beta} \in S'_{\boldsymbol{\alpha}}$ with $\mathtt{Hp}(f,[\boldsymbol{z}],x_i)(\boldsymbol{\beta}) \neq 0$, there exists an open connected set $S''_{\boldsymbol{\alpha}} \subseteq S'_{\boldsymbol{\alpha}}$ and f is open delineable on $S''_{\boldsymbol{\alpha}}$ w.r.t. $\overline{\mathtt{Hp}}(f,i)$ and $\widetilde{\mathtt{Hp}}(f,i)$. From union property of open delineable (Proposition 27), f is open delineable on $S'_{\boldsymbol{\alpha}}$ w.r.t. $\overline{\mathtt{Hp}}(f,i)$ and $\widetilde{\mathtt{Hp}}(f,i)$.

To summarize, the above discussion shows that for any point $\boldsymbol{\alpha} \in S$, there exists an open connected set $S_{\boldsymbol{\alpha}} \subseteq S$ such that $\boldsymbol{\alpha} \in S_{\boldsymbol{\alpha}}$ and f is open delineable on $S_{\boldsymbol{\alpha}}$ w.r.t. $\overline{\mathtt{Hp}}(f,i)$ and $\widetilde{\mathtt{Hp}}(f,i)$. By the nonempty intersection property of open delineable (Proposition 26) and the fact that S is connected, $f(\boldsymbol{x}_n)$ is open delineable on S w.r.t. $\overline{\mathtt{Hp}}(f,i)$ and $\widetilde{\mathtt{Hp}}(f,i)$ as desired.

Therefore, the theorem is proved by induction. The last statement of the theorem follows from the fact $\mathrm{Zero}(\{\mathtt{Hp}(f,[x_n,\ldots,x_j],x_t) \mid t = j,\ldots,n\}) = \mathrm{Zero}(H_{j-1}(f))$. □

5. Application: Open Sample

As a first application of Theorem 32, we show how to compute open sample based on Algorithm 1.

Definition 33. A *reduced open CAD* of $f(\boldsymbol{x}_n)$ w.r.t. $[x_n,\ldots,x_{j+1}]$ is a set of sample points in $\mathbb{R}^n$ obtained through Algorithm 4.

Algorithm 4. `ReducedOpenCAD`

```
Input: $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$ and an open sample $S_{\texttt{Hp}(f,[x_n,\ldots,x_{j+1}]),\texttt{Hp}(f,[x_n,\ldots,x_{j+1}],x_{j+1})}$ in $\mathbb{R}^j$
Output: A set of sample points in $\mathbb{R}^n$
 1: $O := S_{\texttt{Hp}(f,[x_n,\ldots,x_{j+1}]),\texttt{Hp}(f,[x_n,\ldots,x_{j+1}],x_{j+1})}$
 2: for $i$ from $j+2$ to $n+1$ do
 3:     $P := \emptyset$
 4:     for $\boldsymbol{\alpha}$ in $O$ do
 5:         if $i \le n$ then
 6:             $P := P \bigcup (\boldsymbol{\alpha} \boxplus \texttt{SPOne}(\texttt{Hp}(f,[x_n,\ldots,x_i])(\boldsymbol{\alpha},x_{i-1}), \texttt{Hp}(f,[x_n,\ldots,x_i],x_i)(\boldsymbol{\alpha},x_{i-1})))$
 7:         else
 8:             $P := P \bigcup (\boldsymbol{\alpha} \boxplus \texttt{SPOne}(f(\boldsymbol{\alpha},x_n), f(\boldsymbol{\alpha},x_n)))$
 9:         end if
10:     end for
11:     $O := P$
12: end for
13: return $O$
```

The following corollary of Theorem 32 shows that the reduced open CAD owns the property of open delineability.

Corollary 34. *A reduced open CAD of $f(\boldsymbol{x}_n)$ w.r.t. $[x_n,\ldots,x_{j+1}]$ is an open sample defined by $f(\boldsymbol{x}_n)$.*

Example 7. We illustrate the main steps of Algorithm 4. using the polynomial f from Examples 3 and 5.

In: $f = (x_3^2 + x_2^2 + x_1^2 - 1)(4x_3 + 3x_2 + 2x_1 - 1) \in \mathbb{Z}[x_1, x_2, x_3]$
$S_{\texttt{Hp}(f,[x_3,x_2]),\texttt{Hp}(f,[x_3,x_2],x_2)} = \{-2, -\frac{27}{32}, 0, \frac{63}{64}, 2\}$ in $\mathbb{R}$.
1: $O := \{-2, -\frac{27}{32}, 0, \frac{63}{64}, 2\}$ (O has 5 elements, $\boldsymbol{\alpha}_1, \ldots, \boldsymbol{\alpha}_5$)
3: $P := \emptyset$
6: $P := P \bigcup (\boldsymbol{\alpha}_1 \boxplus \texttt{SPOne}(\texttt{Hp}(f,[x_3])(\boldsymbol{\alpha}_1,x_2), \texttt{Hp}(f,[x_3],x_3)(\boldsymbol{\alpha}_1,x_2)))$
$P := P \bigcup (\boldsymbol{\alpha}_2 \boxplus \texttt{SPOne}(\texttt{Hp}(f,[x_3])(\boldsymbol{\alpha}_2,x_2), \texttt{Hp}(f,[x_3],x_3)(\boldsymbol{\alpha}_2,x_2)))$
$\vdots$
$P := P \bigcup (\boldsymbol{\alpha}_5 \boxplus \texttt{SPOne}(\texttt{Hp}(f,[x_3])(\boldsymbol{\alpha}_5,x_2), \texttt{Hp}(f,[x_3],x_3)(\boldsymbol{\alpha}_5,x_2)))$
11 $O := P$ (O now has 13 elements, $\boldsymbol{\alpha}_1, \ldots, \boldsymbol{\alpha}_{13}$)
3: $P := \emptyset$
8: $P := P \bigcup (\boldsymbol{\alpha}_1 \boxplus \texttt{SPOne}(\texttt{Hp}(f)(\boldsymbol{\alpha}_1,x_3), \texttt{Hp}(f)(\boldsymbol{\alpha}_1,x_3)))$
$P := P \bigcup (\boldsymbol{\alpha}_2 \boxplus \texttt{SPOne}(\texttt{Hp}(f)(\boldsymbol{\alpha}_2,x_3), \texttt{Hp}(f)(\boldsymbol{\alpha}_2,x_3)))$
$\vdots$
$P := P \bigcup (\boldsymbol{\alpha}_{13} \boxplus \texttt{SPOne}(\texttt{Hp}(f)(\boldsymbol{\alpha}_{13},x_3), \texttt{Hp}(f)(\boldsymbol{\alpha}_{13},x_3)))$
11: $O := P$ (O now has 36 elements, $\boldsymbol{\alpha}_1, \ldots, \boldsymbol{\alpha}_{36}$)

Out O

Remark 35. As an application of Theorem 32, we could design a CAD-like method to get an open sample defined by $f(\boldsymbol{x}_n)$ for a given polynomial $f(\boldsymbol{x}_n)$. Roughly speaking, if we have already got an open sample defined by $\texttt{Hp}(f, [x_n, \ldots, x_j])$ in $\mathbb{R}^{j-1}$, according to Theorem 32, we could obtain an open sample defined by f in $\mathbb{R}^n$. That process could be done recursively.

In the definition of Hp, we first choose m variables from $\{x_1, ..., x_n\}$, compute all projection polynomials under all possible orders of those m variables, and then compute the gcd of all those projection polynomials. Therefore, Theorem 32 provides us many ways for designing various algorithms for computing open samples. For example, we may set $m = 2$ and choose $[x_n, x_{n-1}]$, $[x_{n-2}, x_{n-3}]$, etc. successively in each step. Because there are only two different orders for two variables, we compute the gcd of two projection polynomials under the two orders in each step. Algorithm 5 is based on this choice.

Algorithm 5. `HpTwo`

Input: A polynomial $f \in \mathbb{Z}[\boldsymbol{x}_n]$ of level n.

Output: An open sample defined by f, *i.e.*, a set of sample points which contains at least one point from each connected component of $f \neq 0$ in $\mathbb{R}^n$

```
$g := f$; $L_1 := \{\}$; $L_2 := \{\}$;
while $i \geq 3$ do
    $L_1 := L_1 \bigcup \overline{\texttt{Hp}}(g, i-1)$;
    $L_2 := L_2 \bigcup \widetilde{\texttt{Hp}}(g, i-1)$;
    $g := \texttt{Hp}(g, [x_i, x_{i-1}])$;
    $i := i - 2$;
end while
if $i = 2$ then
    $L_1 := L_1 \bigcup \overline{\texttt{Hp}}(g, i)$;
    $L_2 := L_2 \bigcup \widetilde{\texttt{Hp}}(g, i)$;
    $g := \texttt{Hp}(g, [x_i])$;
end if
$S$:=SPOne($L_1^{[1]}, L_2^{[1]}$);
$C$:= OpenSP($L_1, L_2, S$);
return $C$.
```

Remark 36. If $\texttt{Hp}(f, [x_n, x_{n-1})] \neq \texttt{Bp}(f, [x_n, x_{n-1}])$ and $n > 3$, it is obvious that the scale of projection in Algorithm 5 is smaller than that of open CAD in Definition 14.

Remark 37. It should be mentioned that there are some non-CAD methods for computing sample points in semi-algebraic sets, such as critical point method. For related work, see for example, Basu et al. (1998); Safey El Din and Schost (2003); Hong and Safey El Din (2012). Mohab Safey el Din developed a Maple package `RAGlib`[1] based on their work, which can test semi-definiteness of polynomials.

---

[1] http://www-polsys.lip6.fr/~safey/RAGLib/distrib.html

6. Application: Polynomial Inequality Proving

In this section, we combined the idea of `Hp` and the simplified CAD projection operator `Np` we introduced previously in Han et al. (2013), to get a new algorithm for testing semi-definiteness of polynomials.

Definition 38. (Han et al., 2013) Suppose $f \in \mathbb{Z}[\boldsymbol{x}_n]$ is a polynomial of level n. Define

$$\begin{aligned}
&\mathrm{Oc}(f, x_n) = \mathrm{sqrfree}_1(\mathrm{lc}(f, x_n)), \mathrm{Od}(f, x_n) = \mathrm{sqrfree}_1(\mathrm{discrim}(f, x_n)),\\
&\mathrm{Ec}(f, x_n) = \mathrm{sqrfree}_2(\mathrm{lc}(f, x_n)), \mathrm{Ed}(f, x_n) = \mathrm{sqrfree}_2(\mathrm{discrim}(f, x_n)),\\
&\mathrm{Ocd}(f, x_n) = \mathrm{Oc}(f, x_n) \cup \mathrm{Od}(f, x_n),\\
&\mathrm{Ecd}(f, x_n) = \mathrm{Ec}(f, x_n) \cup \mathrm{Ed}(f, x_n).
\end{aligned}$$

The *secondary* and *principal parts* of the projection operator `Np` are defined as

$$\begin{aligned}
\mathtt{Np}_1(f, [x_n]) =& \mathrm{Ocd}(f, x_n),\\
\mathtt{Np}_2(f, [x_n]) =& \{\prod_{g \in \mathrm{Ecd}(f, x_n) \setminus \mathrm{Ocd}(f, x_n)} g\}.
\end{aligned}$$

If L is a set of polynomials of level n, define

$$\begin{aligned}
\mathtt{Np}_1(L, [x_n]) &= \bigcup_{g \in L} \mathrm{Ocd}(g, x_n),\\
\mathtt{Np}_2(L, [x_n]) &= \bigcup_{g \in L} \{\prod_{h \in \mathrm{Ecd}(g, x_n) \setminus \mathtt{Np}_1(L, [x_n])} h\}.
\end{aligned}$$

Based on the projection operator `Np`, we proposed an algorithm, `DPS`, in (Han et al., 2013) for testing semi-definiteness of polynomials. Algorithm `DPS` takes a polynomial $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$ as input, and returns whether or not $f(\boldsymbol{x}_n) \geq 0$ on $\mathbb{R}^n$. The readers are referred to (Han et al., 2013) for the details of `DPS`.

The projection operator `Np` is extended and defined in the next definition.

Definition 39. Let $f \in \mathbb{Z}[x_1, \ldots, x_n]$ with level n. Denote $[\boldsymbol{y}] = [y_1, \ldots, y_m]$, for $1 \leq m \leq n$, where $y_i \in \{x_1, \ldots, x_n\}$ for $1 \leq i \leq m$ and $y_i \neq y_j$ for $i \neq j$. Define

$$\mathtt{Np}(f, [x_i]) = \mathtt{Np}_2(f, [x_i]), \quad \mathtt{Np}(f, [x_i], x_i) = \prod_{g \in \mathtt{Np}_1(f, [x_i])} g.$$

For $m (m \geq 2)$ and $i (1 \leq i \leq m)$, $\mathtt{Np}(f, [\boldsymbol{y}], y_i)$ and $\mathtt{Np}(f, [\boldsymbol{y}])$ are defined recursively as follows.

$$\begin{aligned}
&\mathtt{Np}(f, [\boldsymbol{y}], y_i) = \mathtt{Bp}(\mathtt{Np}(f, [\hat{\boldsymbol{y}}]_i), y_i),\\
&\mathtt{Np}(f, [\boldsymbol{y}]) = \gcd(\mathtt{Np}(f, [\boldsymbol{y}], y_1), \ldots, \mathtt{Np}(f, [\boldsymbol{y}], y_m)),
\end{aligned}$$

where $[\hat{\boldsymbol{y}}]_i = [y_1, \ldots, y_{i-1}, y_{i+1}, \ldots, y_m]$. Define

$$\overline{\mathtt{Np}}(f, i) = \{f, \mathtt{Np}(f, [x_n]), \ldots, \mathtt{Np}(f, [x_n, \ldots, x_i)]\},$$

and

$$\widetilde{\mathtt{Np}}(f, i) = \{f, \mathtt{Np}(f, [x_n], x_n), \ldots, \mathtt{Np}(f, [x_n, \ldots, x_i], x_i)\}.$$

Theorem 40. *(Han et al., 2013) Given a positive integer* $n \geq 2$. *Let* $f \in \mathbb{Z}[\boldsymbol{x}_n]$ *be a non-zero squarefree polynomial and* U *a connected component of* $\mathtt{Np}(f,[x_n]) \neq 0$ *in* $\mathbb{R}^{n-1}$. *If the polynomials in* $\mathtt{Np}_1(f,[x_n])$ *are semi-definite on* U, *then* f *is delineable on* $V = U \setminus \bigcup_{h \in \mathtt{Np}_1(f,[x_n])} \mathrm{Zero}(h)$.

Lemma 41. *(Han et al., 2013) Given a positive integer* $n \geq 2$. *Let* $f \in \mathbb{Z}[\boldsymbol{x}_n]$ *be a squarefree polynomial with level* n *and* U *a connected open set of* $\mathtt{Np}(f,[x_n]) \neq 0$ *in* $\mathbb{R}^{n-1}$. *If* $f(\boldsymbol{x}_n)$ *is semi-definite on* $U \times \mathbb{R}$, *then the polynomials in* $\mathtt{Np}_1(f,[x_n])$ *are all semi-definite on* U.

Now, we can rewrite Theorem 40 in another way.

Proposition 42. *Let* $f \in \mathbb{Z}[\boldsymbol{x}_n]$ *be a squarefree polynomial with level* n *and* U *a connected component of* $\mathtt{Np}(f,[x_n]) \neq 0$ *in* $\mathbb{R}^{n-1}$. *If the polynomials in* $\mathtt{Np}_1(f,[x_n])$ *are semi-definite on* U, *then* f *is open delineable on* U *w.r.t.* $\overline{\mathtt{Np}}(f,n)$ *and* $\widetilde{\mathtt{Np}}(f,n)$.

Theorem 43. *Let* j *be an integer and* $2 \leq j \leq n$. *For any given polynomial* $f(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$, *and any open connected set* U *of* $\mathtt{Np}(f,[x_n,\ldots,x_j]) \neq 0$ *in* $\mathbb{R}^{j-1}$, *let*

$$S = U \setminus \mathrm{Zero}(\{\mathtt{Np}(f,[x_n,\ldots,x_j],x_t) \mid t = j,\ldots,n\}).$$

If the polynomials in $\bigcup_{i=0}^{n-j} \mathtt{Np}_1(f,[x_{n-i}])$ *are all semi-definite on* $U \times \mathbb{R}^{n-j}$, $f(\boldsymbol{x}_n)$ *is open delineable on* S *w.r.t.* $\overline{\mathtt{Np}}(f,j)$ *and* $\widetilde{\mathtt{Np}}(f,j)$.

Proof. Notice that the proof of Theorem 32 only uses the properties of open delineable (Propositions 24-27) and Proposition 31(f is open delineable on every open connected set defined by $\mathtt{Bp}(f,[x_n]) \neq 0$ in $\mathbb{R}^{n-1}$ w.r.t. $\overline{\mathtt{Hp}}(f,n)$ and $\widetilde{\mathtt{Hp}}(f,n)$). According to Proposition 42, f is open delineable on every open connected set defined by $\mathtt{Np}(f,[x_n]) \neq 0$ in $\mathbb{R}^{n-1}$ w.r.t. $\overline{\mathtt{Np}}(f,n)$ and $\widetilde{\mathtt{Np}}(f,n)$. The same proof of Theorem 32 will yield the conclusion of the theorem. □

Theorem 43 and Proposition 41 provide us a new way to decide the non-negativity of a polynomial as stated in the next theorem.

Theorem 44. *Let* $f \in \mathbb{Z}[\boldsymbol{x}_n]$ *be a squarefree polynomial with level* n *and* U *a connected open set of* $\mathtt{Np}(f,[x_n,\ldots,x_j]) \neq 0$ *in* $\mathbb{R}^{j-1}$. *Denote* $S = U \setminus \mathrm{Zero}(\{\mathtt{Np}(f,[x_n,\ldots,x_j],x_t) \mid t = j,\ldots,n\})$. *The necessary and sufficient condition for* $f(\boldsymbol{x}_n)$ *to be positive semi-definite on* $U \times \mathbb{R}^{n-j+1}$ *is the following two conditions hold.*
(1) *The polynomials in* $\bigcup_{i=0}^{n-j} \mathtt{Np}_1(f,[x_{n-i}])$ *are all semi-definite on* $U \times \mathbb{R}^{n-j}$.
(2) *There exists a point* $\boldsymbol{\alpha} \in S$ *such that* $f(\boldsymbol{\alpha},x_j,\ldots,x_n)$ *is positive semi-definite on* $\mathbb{R}^{n-j+1}$.

Based on the above theorems, it is easy to design some different algorithms (depending on the choice of j) to prove polynomial inequality. For example, the algorithm `PSD-HpTwo` for deciding whether a polynomial is positive semi-definite, which we will introduce later, is based on Theorem 44 when $j = n-1$ (Proposition 45).

Proposition 45. *Given a positive integer* $n \geq 3$*. Let* $f \in \mathbb{Z}[\boldsymbol{x}_n]$ *be a squarefree polynomial with level* n *and* U *a connected open set of* $\mathtt{Np}(f, [x_n, x_{n-1}]) \neq 0$ *in* $\mathbb{R}^{n-2}$*. Denote* $S = U \backslash \mathrm{Zero}(\mathtt{Np}(f, [x_n, x_{n-1}], x_n), \mathtt{Np}(f, [x_n, x_{n-1}], x_{n-1}))$*.*
The necessary and sufficient condition for $f(\boldsymbol{x}_n)$ *to be positive semi-definite on* $U \times \mathbb{R}^2$ *is the following two conditions hold.*
(1) *The polynomials in either* $\mathtt{Np}_1(f, [x_n])$ *or* $\mathtt{Np}_1(f, [x_{n-1}])$ *are semi-definite on* $U \times \mathbb{R}$*.*
(2) *There exists a point* $\boldsymbol{\alpha} \in S$ *such that* $f(\boldsymbol{\alpha}, x_{n-1}, x_n)$ *is positive semi-definite on* $\mathbb{R}^2$*.*

Algorithm 6. `PSD-HpTwo`
Input: An irreducible polynomial $f \in \mathbb{Z}[\boldsymbol{x}_n]$.
Output: Whether or not $\forall \boldsymbol{\alpha}_n \in \mathbb{R}^n$, $f(\boldsymbol{\alpha}_n) \geq 0$.

```
if $n \leq 2$ then
    if DPS($f(x_n)$)=false then return false
    end if
else
    $L_1 := \mathtt{Np}_1(f, [x_n]) \bigcup \mathtt{Np}_1(f, [x_{n-1}])$
    $L_2 := \mathtt{Np}(f, [x_n, x_{n-1}])$
    for $g$ in $L_1$ do
        if PSD-HpTwo($g$) =false then return false
        end if
    end for
    $C_{n-2}$ := A reduced open CAD of $L_2$ w.r.t. $[x_{n-2}, \ldots, x_2]$, which satisfies that
      Zero($\mathtt{Np}(f, [x_n, x_{n-1}], x_n), \mathtt{Np}(f, [x_n, x_{n-1}], x_{n-1})$) $\cap C_{n-2} = \emptyset$.
    if $\exists \boldsymbol{\alpha}_{n-2} \in C_{n-2}$ such that DPS($f(\boldsymbol{\alpha}_{n-2}, x_{n-1}, x_n)$)=false then return false
    end if
end if
return true
```

7. Application: Copositive problem

Definition 46. A real $n \times n$ matrix A_n is said to be *copositive* if $\boldsymbol{x}_n A_n \boldsymbol{x}_n^T \geq 0$ for every nonnegative vector $\boldsymbol{x}_n$. For convenience, we also say the form $\boldsymbol{x}_n A_n \boldsymbol{x}_n^T$ is copositive if A_n is copositive.

The collection of all copositive matrices is a proper cone; it includes as a subset the collection of real positive-definite matrices. For example, xy is copositive but it is not positive semi-definite.

In general, to check whether a given integer square matrix is not copositive, is NP-complete (Murty and Kabadi, 1987). This means that every algorithm that solves the problem, in the worst case, will require at least an exponential number of operations, unless P=NP. For that reason, it is still valuable for the existence of so many incomplete algorithms discussing some special kinds of matrices (Parrilo, 2000). For small values of n (≤ 6), some necessary and sufficient conditions have been constructed (Andersson et al., 1995; Hadeler, 1983). We refer the reader to (Hiriart-Urruty and Seeger, 2010) for a more detailed introduction to copositive matrices.

From another viewpoint, this is a typical real quantifier elimination problem, which can be solved by standard tools of real quantifier elimination (*e.g.*, using typical CAD). Thus, any CAD based QE algorithm can serve as a complete algorithm for deciding copositive matrices theoretically. Unfortunately, such algorithm is not efficient in practice for the CAD algorithm is of doubly exponential time complexity.

To test the copositivity of the form $\boldsymbol{x}_n A_n \boldsymbol{x}_n^T$, is equivalent to test the nonnegativity of the form $(x_1^2, \dots, x_n^2) A_n (x_1^2, \dots, x_n^2)^T$. In this section, we give a singly exponential incomplete algorithm with time complexity $\mathcal{O}(n^2 4^n)$ based on the new projection operator proposed in the last section and Theorem 44.

Let us take an example to illustrate our idea. Let

$$F := ax^4 + bx^2y^2 + cy^4 + dx^2 + ey^2 + f,$$

be a squarefree polynomial, where $a, b, c \in \mathbb{Z}, d, e, f \in \mathbb{Z}[\boldsymbol{z}_n]$ and $a \neq 0,\ c \neq 0$.

To test the nonnegativity of F, we could apply typical CAD-based methods directly, *i.e.*, we can use Brown's projection operator. In general, we have

$$\mathtt{Bp}(F, [x]) = (cy^4 + ey^2 + f)a(4acy^4 + 4aey^2 + 4af - b^2y^4 - 2by^2d - d^2).$$

We then eliminate y,

$$\begin{aligned} &\mathtt{Bp}(F, [x, y]) = \mathtt{Bp}(\mathtt{Bp}(F, [x]), y) \\ =&fac(4fc - e^2)(d^2c - edb + fb^2)(4af - d^2)(4ac - b^2)(4afc - ae^2 - d^2c + edb - fb^2). \end{aligned}$$

If d, e are polynomials of degree 2 and f is a polynomial of degree 4 (copositive problem is in this case), the degree of the polynomial $\mathtt{Bp}(F, [x, y])$ is 20 while the original problem is of degree 4 only. That could help us understand why typical CAD-based methods do not work for copositive problems with more than 5 variables in practice.

Now, we apply our new projection operator. Notice that

$$\mathrm{Res}(F, \frac{\partial F}{\partial x}, x) = 16(cy^4 + ey^2 + f)F_1^2,$$

where $F_1 = a(4acy^4 + 4aey^2 + 4af - b^2y^4 - 2by^2d - d^2)$.

If $cy^4 + ey^2 + f$ and F_1 are nonzero and squarefree, $\mathtt{Np}(F, [x]) = F_1$. Thus, in order to test the nonnegativity of F, we only need to test the semi-definitness of $cy^4 + ey^2 + f$, choose sample points defined by $\mathtt{Np}(F, [x]) \neq 0$ (we also require that cy^4+ey^2+f does not vanish at those sample points) and test the nonnegativity of F at these sample points.

On the other side,

$$\mathrm{Res}(F, \frac{\partial F}{\partial y}, y) = 16(ax^4 + dx^2 + f)F_2^2,$$

where $F_2 = c(4cx^4a + 4cdx^2 + 4fc - b^2x^4 - 2bx^2e - e^2)$.

Similarly, if ax^4+dx^2+f and F_2 are nonzero and squarefree, $\mathtt{Np}(F, [y]) = F_2$. In order to test the nonnegativity of F, we only need to test the semi-definitness of ax^4+dx^2+f, choose sample points defined by $\mathtt{Np}(F, [y]) \neq 0$ (we also require that ax^4+dx^2+f does not vanish at those sample points) and test the nonnegativity of F at these sample points.

Under some "generic" conditions (*i.e.*, some polynomials are nonzero and squarefree), we only need to test the semi-definitness of $ax^4 + dx^2 + f$ and $cy^4 + ey^2 + f$, choose sample points T_2 defined by $\mathtt{Np}(F, [x, y]) = \gcd(\mathtt{Bp}(\mathtt{Np}(F, [x]), y),\ \mathtt{Bp}(\mathtt{Np}(F, [y]), x)) = (4ac - b^2)(4afc - ae^2 - d^2c + edb - fb^2) \neq 0$ (we also require that $\mathrm{Res}(\mathtt{Np}(F, [x]), y)$ does not vanish at T_2), obtain sample points T_1 defined by $\mathtt{Np}(F,[x]) \neq 0$ from T_2 (we also require that cy^4+ey^2+f does not vanish at T_1) and test the nonnegativity of F at T_1.

Again, if d, e are polynomials of degree 2 and f is a polynomial of degree 4, both the degree of $\mathtt{Np}(F,[x])$ and $\mathtt{Np}(F,[x,y])$ are exactly 4. It indicates that our new projection operator may control the degrees of polynomials in projection sets. Moreover, we point out that

$$\begin{aligned}\mathtt{Np}(F,[x,y]) &= (4afc-ae^2-d^2c+edb-fb^2)\\ &= 4\det\left(\begin{bmatrix} a & \frac{b}{2} & \frac{d}{2} \\ \frac{b}{2} & c & \frac{e}{2} \\ \frac{d}{2} & \frac{e}{2} & f \end{bmatrix}\right).\end{aligned}$$

Before giving the result, we introduce some new notations and lemmas for convenience.

Definition 47 (Sub-sequence)**.** An array I is called a *sub-sequence* of sequence $\{1,\dots,n\}$ if for any i-th component of I, $I[i] \in \{1,\dots,n\}$ and $I[i] < I[i+1]$ for $i=1,\dots,|I|-1$.

For a sub-sequence I of $\{1,\dots,n\}$ with $m=|I|$, denote $\boldsymbol{x}_I = [x_{I[1]},\dots,x_{I[m]}]$, $\overline{\boldsymbol{x}_I} = \{x_i \mid i \notin I\}$ and $A_I = (a_{I[i],I[j]})_{1\le i,j\le m}$ the sub-matrix of A_n.

Let

$$f(\boldsymbol{x}_n) = \sum_{1\le i,j\le n} a_{i,j}x_ix_j + \sum_{i=1}^{n}(a_{i,n+1}+a_{n+1,i})x_i + a_{n+1,n+1} = (\boldsymbol{x}_n,1)A_{n+1}(\boldsymbol{x}_n,1)^T \quad (3)$$

be a quadratic polynomial in $\boldsymbol{x}_n$, where $A_{n+1} = (a_{i,j})_{(n+1)\times(n+1)}$, $a_{i,j}=a_{j,i} \in \mathbb{Z}[\boldsymbol{z}_s]$ for $1 \le i,j \le n+1$. Set $F(\boldsymbol{x}_n) = f(x_1^2,\dots,x_n^2)$. It is not hard to see that (please refer to the proof of Theorem 49), for a given sub-sequence I of $\{1,\dots,n\}$ with length m, there exist some polynomials $p_1,\dots,p_{m+1} \in \mathbb{Z}[\boldsymbol{z}_s,\overline{\boldsymbol{x}_I}]$ such that $F(\boldsymbol{x}_n) = (\boldsymbol{x}_I^2,1)P_I(\boldsymbol{x}_I^2,1)^T$ where

$$P_I = \begin{bmatrix} A_I & (p_1,\cdots,p_m)^T \\ (p_1,\cdots,p_m) & p_{m+1} \end{bmatrix}.$$

For convenience, we denote $P_{[1,\dots,m]}$ by $P_{m+1}(x_{m+1},\dots,x_n)$ or simply P_{m+1}. In particular, $P_{n+1} = A_{n+1}$ and $F(\boldsymbol{x}_n) = (x_1^2,\dots,x_i^2,1)P_{i+1}(x_1^2,\dots,x_i^2,1)^T$.

Example 8. Suppose $F(x_1,x_2,x_3) = x_1^4+2x_2^4+4x_1^2x_2^2-2x_1^2x_3^2+4x_2^2x_3^2+8x_1^2z^2+5x_3^4+z^4$. If $I=[1,2]$, then $\boldsymbol{x}_I = [x_1,x_2], \overline{\boldsymbol{x}_I} = \{x_3\}$,

$$P_I = \begin{bmatrix} 1 & 2 & 4z^2-x_3^2 \\ 2 & 2 & 2x_3^2 \\ 4z^2-x_3^2 & 2x_3^2 & 5x_3^4+z^4 \end{bmatrix}.$$

If $I=[1]$, then $\boldsymbol{x}_I = [x_1], \overline{\boldsymbol{x}_I} = \{x_2,x_3\}$,

$$P_I = \begin{bmatrix} 1 & 2x_2^2+4z^2-x_3^2 \\ 2x_2^2+4z^2-x_3^2 & 2x_2^4+5x_3^4+z^4 \end{bmatrix}.$$

Lemma 48. *Suppose R is a square matrix with order n, P is an invertible square matrix with order $k < n$, $Q \in \mathbb{R}^{k\times(n-k)}$, $M \in \mathbb{R}^{(n-k)\times k}$ and $N \in \mathbb{R}^{(n-k)\times(n-k)}$. If R can be written as partitioned matrix*

$$R = \begin{bmatrix} P & Q \\ M & N \end{bmatrix},$$

then

$$\det(R) = \det(P)\det(N - MP^{-1}Q).$$

Proof. It is a well known result in linear algebra. $\square$

For a square matrix M, we use $M^{(i,j)}$ to denote the determinant of the sub-matrix obtained by deleting the i-th row and the j-th column of M.

Theorem 49. *Suppose $f \in \mathbb{Z}[\boldsymbol{z}_s][\boldsymbol{x}_n]$ is defined as in (3). Set $F(\boldsymbol{x}_n) = f(x_1^2, \ldots, x_n^2)$. If*

(1) $F(\boldsymbol{x}_n)|_{x_i=0}$ is nonzero and squarefree for any $i \in \{1, \ldots, n\}$;

(2) $\det(A_I) = P_I^{(|I|+1,|I|+1)}$ is nonzero and squarefree for any sub-sequence I of $\{1, \ldots, n\}$, and $\gcd(A_I^{(1,1)}, \ldots, A_I^{(|I|,|I|)}) = 1$ for any any sub-sequence $|I| \geq 2$ of $\{1, \ldots, n\}$;

(3) $\gcd(P_I^{(1,1)}, \ldots, P_I^{(|I|,|I|)}) = 1$ for any sub-sequence $|I| \geq 2$ of $\{1, \ldots, n\}$;

(4) $\det(P_I)$ is nonzero and squarefree for any sub-sequence I of $\{1, \ldots, n\}$;

(5) $\gcd(\det(P_I), \det(A_I)) = 1$ for any sub-sequence I of $\{1, \ldots, n\}$,

then $\mathtt{Np}(F, [\boldsymbol{x}_n]) = \det(A_n)\det(A_{n+1})$.

Proof. We prove the theorem by induction on n.

If $n = 1$, $F(\boldsymbol{x}_1) = a_{1,1}x_1^4 + 2a_{1,2}x_1^2 + a_{2,2}$. Then

$$\mathrm{Res}(F, \frac{\partial F}{\partial x_1}, x_1) = 256a_{1,1}^2 a_{2,2}(a_{1,1}a_{2,2} - a_{1,2}^2)^2.$$

By conditions (1), (2), (4) and (5), $a_{1,1} \neq 0, a_{2,2} \neq 0, a_{1,1}a_{2,2} - a_{1,2}^2 \neq 0$, and $a_{2,2}$ and $a_{1,1}a_{2,2} - a_{1,2}^2$ are two coprime squarefree polynomials. Thus, $\mathtt{Np}(F, [\boldsymbol{x}_1]) = a_{1,1}(a_{1,1}a_{2,2} - a_{1,2}^2) = \det(A_1)\det(A_2)$.

Assume that the conclusion holds for any quadratic polynomials with k variables where $1 \leq k < n$. When $n = k$, let I be a sub-sequence of $\{1, \ldots, n\}$ with $|I| = n - 1$. Without loss of generality, we assume that $I = [1, \ldots, n-1]$. Set $A_I = (a_{i,j})_{1\leq i,j<n}$, $B = (a_{1,n}, \ldots, a_{n-1,n})$, $C = (a_{1,n+1}, \ldots, a_{n-1,n+1})$, and $D = a_{n,n}x_n^4 + 2a_{n,n+1}x_n^2 + a_{n+1,n+1}$. Then, $F(\boldsymbol{x}_n)$ could be written as

$$\begin{aligned} F(\boldsymbol{x}_n) = &\sum_{1\leq i,j<n} a_{i,j}x_i^2x_j^2 + \sum_{i=1}^{n-1}(a_{i,n} + a_{n,i})x_i^2x_n^2 + (a_{n,n}x_n^4 + 2a_{n,n+1}x_n^2 + a_{n+1,n+1}) \\ =&(x_1^2, \ldots, x_{n-1}^2, 1)P_I(x_1^2, \ldots, x_{n-1}^2, 1)^T, \end{aligned}$$

where

$$P_I = \begin{bmatrix} A_I & (Bx_n^2 + C)^T \\ Bx_n^2 + C & D \end{bmatrix}.$$

By assumption, $F|_{x_i=0}$ is squarefree for $i \in I$, and $\det(A_I')$ is nonzero and squarefree, $\gcd(A_{I'}^{(1,1)}, \ldots, A_{I'}^{(|I'|,|I'|)}) = 1$, $\gcd(P_{I'}^{(1,1)}, \ldots, P_{I'}^{(|I'|,|I'|)}) = 1$, $\gcd(\det(P_{I'}), \det(A_{I'})) = 1$

for any sub-sequence I' of I with $|I'| \leq n-1$. Thus, by induction hypothesis, $\mathtt{Np}(F(\boldsymbol{x}_n), [\boldsymbol{x}_I]) = \det(A_I)\det(P_I)$.

In the following, we compute $\det(P_I)$. By assumption, $\det(A_I) = P_I^{(n,n)}$ is nonzero and squarefree. According to Lemma 48,

$$\begin{aligned}
\det(P_I) &= \det(A_I)(D - (Bx_n^2 + C)A_I^{-1}(Bx_n^2 + C)^T) \\
&= \det(A_I)\left((a_{n,n} - BA_I^{-1}B^T)x_n^4 + 2(a_{n,n+1} - BA_I^{-1}C^T)x_n^2 + a_{n+1,n+1} - CA_I^{-1}C^T\right) \\
&= \det(A_I)(\lambda x_n^4 + 2\mu x_n^2 + \nu),
\end{aligned}$$

where $\lambda = (a_{n,n} - BA_I^{-1}B^T)$, $\mu = a_{n,n+1} - BA_I^{-1}C^T$, $\nu = a_{n+1,n+1} - CA_I^{-1}C^T$. By Lemma 48 again,

$$\begin{aligned}
\det(A_I)\lambda &= \det(A_I)(a_{n,n} - BA_I^{-1}B^T) \\
&= \det\left(\begin{bmatrix} A_I & B^T \\ B & a_{n,n} \end{bmatrix}\right) \\
&= A_{n+1}^{(n+1,n+1)}, \qquad (4) \\
\det(A_I)\nu &= \det(A_I)(a_{n+1,n+1} - CA_I^{-1}C^T) \\
&= \det\left(\begin{bmatrix} A_I & C^T \\ C & a_{n+1,n+1} \end{bmatrix}\right) \\
&= A_{n+1}^{(n,n)}, \qquad (5)
\end{aligned}$$

Thus, both $\det(A_I)\lambda$ and $\det(A_I)\nu$ are the determinants of some principal sub-matrices of A_{n+1} with order n, respectively.

Let $H = \det(P_I)$, according to Lemma 48, it is clear that

$$\begin{aligned}
\mathrm{Res}(H, \frac{\partial H}{\partial x_n}, x_n) &= 256\det(A_I)^7\lambda^2\nu(\mu^2 - \lambda\nu)^2 \\
&= 256\det(A_I)^7\lambda^2\nu\det\left(\begin{bmatrix} \lambda & \mu \\ \mu & \nu \end{bmatrix}\right)^2 \\
&= 256\det(A_I)^7\lambda^2\nu\det\left(\begin{bmatrix} a_{n,n} & a_{n,n+1} \\ a_{n+1,n} & a_{n+1,n+1} \end{bmatrix} - \begin{pmatrix} B \\ C \end{pmatrix} A_I^{-1} \begin{pmatrix} B \\ C \end{pmatrix}^T\right)^2 \\
&= 256\det(A_I)^5\lambda^2\nu\det\left(\begin{bmatrix} A_I & B^T & C^T \\ B & a_{n,n} & a_{n,n+1} \\ C & a_{n+1,n}, & a_{n+1,n+1} \end{bmatrix}\right)^2 \\
&= 256\det(A_I)^5\lambda^2\nu\det(A_{n+1})^2 \\
&= 256\det(A_I)^2\left(A_{n+1}^{(n+1,n+1)}\right)^2 A_{n+1}^{(n,n)}\det(A_{n+1})^2 \\
&= 256(A_n^{(n,n)})^2\det(A_n)^2 A_{n+1}^{(n,n)}\det(A_{n+1})^2. \qquad (6)
\end{aligned}$$

Since $\det(A_I)$ and $A_{n+1}^{(n+1,n+1)}$ are nonzero, according to (6), we have

$$\mathtt{Np}(F,[\boldsymbol{x}_n],n) = \text{sqrfree}(A_{n+1}^{(n,n)}A_n^{(n,n)}\det(A_n)\det(A_{n+1})).$$

Similarly, for $1 \le i \le n$, we have

$$\mathtt{Np}(F,[\boldsymbol{x}_n],i) = \text{sqrfree}(A_{n+1}^{(i,i)}A_n^{(i,i)}\det(A_n)\det(A_{n+1})). \tag{7}$$

By assumption, $\gcd(A_{n+1}^{(1,1)},\cdots,A_{n+1}^{(n,n)}) = 1$, $\gcd(A_n^{(1,1)},\cdots,A_n^{(n,n)}) = 1$, and $\det(A_n)$, $\det(A_{n+1})$ are two nonzero squarefree polynomials with $\gcd(\det(A_n),\det(A_{n+1})) = 1$, thus

$$\mathtt{Np}(F,[\boldsymbol{x}_n]) = \gcd(\mathtt{Np}(F,[\boldsymbol{x}_n],1),\ldots,\mathtt{Np}(F,[\boldsymbol{x}_n],n)) = \det(A_n)\det(A_{n+1}).$$

That completes the proof. $\square$

Theorem 50. *If the coefficients $a_{i,j}$ of $f(\boldsymbol{x}_n)$ in Theorem 49 are pairwise different real parameters except that $a_{i,j} = a_{j,i}$, and $F(\boldsymbol{x}_n) = f(x_1^2,\ldots,x_n^2)$. Then all the four hypotheses (1)-(5) of Theorem 49 hold. As a result,* $\mathtt{Np}(F,[\boldsymbol{x}_n]) = \det(A_n)\det(A_{n+1})$.

Proof. It is clear that the hypotheses (1) and (2) of Theorem 49 hold. We claim that that for any given m, $|I| = m$, $P_I^{(i,i)}$ and $\det(P_I)$ are pairwise nonconstant different irreducible polynomials in $\mathbb{Z}[\boldsymbol{a}_{i,j}][\boldsymbol{x}_n]$ for all $n \ge m$, so the hypotheses (3),(4) and (5) of Theorem 49 also hold. Here we denote $\boldsymbol{a}_{i,j} = (a_{1,1},\ldots,a_{n+1,n+1})$.

We only prove that $\det(P_I)$ is a nonconstant irreducible polynomial. The other statements of the claim can be proved similarly.

We prove the claim by introduction on n. If $n = m$, it is clear that the claim is true. Assume that the theorem holds for integers $m \le l \le n-1$. We now consider the case $l = n$. Without of loss of generality, we assume that $I = [1,2,\ldots,m]$.

Recall that $F(\boldsymbol{x}_n)$ could be written as

$$F(\boldsymbol{x}_n) = (\boldsymbol{x}_m^2,1)P_{I,F}(\boldsymbol{x}_m^2,1)^T,$$

where

$$P_{I,F} = \begin{bmatrix} A_I & (p_{1,n},\cdots,p_{m,n})^T \\ (p_{1,n},\cdots,p_{m,n}) & p_{m+1,n} \end{bmatrix},$$

and $p_{i,n} = \sum_{j=m+1}^n a_{i,j}x_j^2 + a_{i,n+1}$ for $1 \le i \le m$, and $p_{m+1,n} = \sum_{j=m+1}^n a_{j,j}x_j^4 + \sum_{j=m+1}^n 2a_{n+1,j}x_j^2 + \sum_{m+1\le i<j\le n} 2a_{i,j}x_i^2x_j^2 + a_{n+1,n+1}$. Let $B = (a_{1,n},\ldots,a_{m,n})$, $C = (p_{1,n-1},\ldots,p_{m,n-1})$, and $D = p_{m+1,n} = a_{n,n}x_n^4 + 2a_{n,n+1}x_n^2 + 2\sum_{m+1\le i<n} a_{i,n}x_i^2x_n^2 + a'_{n+1,n+1}$, where $a'_{n+1,n+1}$ is a polynomial with $\deg(a'_{n+1,n+1},x_n) = 0$. Now $P_{I,F}$ can be simply written as

$$P_{I,F} = \begin{bmatrix} A_I & (Bx_n^2+C)^T \\ Bx_n^2+C & D \end{bmatrix}.$$

In the following, we compute $\det(P_{I,F})$. By Lemma 48,

$$\begin{aligned}\det(P_{I,F}) =& \det(A_I)(D-(Bx_n^2+C)A_I^{-1}(Bx_n^2+C)^T)\\ =& \det(A_I)\left((a_{n,n}-BA_I^{-1}B^T)x_n^4 + 2(a_{n,n+1}+\sum_{m+1\le i<n} a_{i,n}x_i^2 - BA_I^{-1}C^T)x_n^2\right.\\ & \left. + a'_{n+1,n+1} - CA_I^{-1}C^T\right)\\ =& \det(A_I)(\lambda x_n^4 + 2\mu x_n^2+\nu),\end{aligned}$$

where $\lambda = (a_{n,n} - BA_I^{-1}B^T)$, $\mu = a_{n,n+1} + \sum_{m+1\le i<n} a_{i,n}x_i^2 - BA_I^{-1}C^T$, $\nu = a'_{n+1,n+1} - CA_I^{-1}C^T$. By Lemma 48,

$$\begin{aligned}\det(A_I)\lambda =& \det(A_I)(a_{n,n}-BA_I^{-1}B^T)\\ =& \det\left(\begin{bmatrix} A_I & B^T\\ B & a_{n,n}\end{bmatrix}\right)\\ \det(A_I)\nu =& \det(A_I)(a'_{n+1,n+1}-CA_I^{-1}C^T)\\ =& \det\left(\begin{bmatrix} A_I & C^T\\ C & a'_{n+1,n+1}\end{bmatrix}\right).\end{aligned}$$

We have $\det(A_I)\lambda = \det(P_{I,G})$, $\det(A_I)\nu = \det(P_{I,H})$, where

$$G(\boldsymbol{x}_m) = (\boldsymbol{x}_m, 1)\left(\begin{bmatrix} A_I & B^T\\ B & a_{n,n}\end{bmatrix}\right)(\boldsymbol{x}_m,1)^T,$$

and

$$H(\boldsymbol{x}_{n-1}) = (\boldsymbol{x}_m, 1)\left(\begin{bmatrix} A_I & C^T\\ C & a'_{n+1,n+1}\end{bmatrix}\right)(\boldsymbol{x}_m,1)^T.$$

By induction, $\det(A_I)\lambda$ and $\det(A_I)\nu$ are two different non-constant irreducible polynomials. Since $\deg(\det(A_I)\mu, a_{n,n+1}) > 0$, and $\deg(\det(A_I)\lambda, a_{n,n+1}) = \deg(\det(A_I)\nu, a_{n,n+1}) = 0$, it is clear that $\det(A_I)\mu \neq \pm(\det(A_I)\lambda\cdot\det(A_I)\nu + 1)$, $\det(A_I)\mu \neq \pm(\det(A_I)\lambda + \det(A_I)\nu)$. Now the result follows from Lemma 51. We are done. $\square$

Lemma 51. *Let $\mathcal{R}$ be a UFD with units ± 1. Let $a,b,c\in\mathcal{R}$, where $b\neq\pm(ac+1)$, $b\neq\pm(a+c)$, and a,c are two non-unit coprime irreducible elements in $\mathcal{R}$, then $T(x) = ax^4+bx^2+c$ is an irreducible polynomial in $\mathcal{R}[x]$.*

Proof. Otherwise, we may assume $T(x) = g(x)h(x)$, where g,h are two nonconstant polynomials in $\mathcal{R}[x]$. Notice that if $\alpha\in\mathcal{R}$ is a root of $T(x)$, then $-\alpha$ is also a root of $T(x)$, thus $(x^2-\alpha^2)$ is a factor of T. Thus, we may assume that $\deg(g)=\deg(h)=2$. Let $g = g_0+g_1x+g_2x^2$, $h = h_0+h_1x+h_2x^2$, where $g_i,h_i\in\mathcal{R}$. By comparing the coefficients of T with gh, we have $c = g_0h_0$, $0 = g_0h_1+g_1h_0$, $h_1g_2+h_2g_1 = 0$. Assume that $c|g_0$, then $c\nmid h_0$. if $h_1\neq 0$, let l be the largest integer such that $c^l|h_1$, then $l+1$ is the largest integer such that $c^{l+1}|g_1$. But $h_1g_2+h_2g_1=0$, so $c|g_2$, and $c|gh$, which contradicts with $(a,c)=1$. We must have $h_1 = 0$, and $g_1 = 0$. We assume that $g_0 = c, h_0 = 1$. Now, there are four cases ($g_2 = \pm a, h_2 = \mp 1$) or ($g_2 = \pm 1, h_2 = \mp a$). All the four cases will contradict with the assumption that $b \neq \pm(ac+1)$, $b \neq \pm(a+c)$. $\square$

Theorem 52. *Suppose* $g(\boldsymbol{x}_n) = \sum_{1\leq i,j\leq n} a_{i,j}x_ix_j = \boldsymbol{x}_n A_n \boldsymbol{x}_n^T$ *is a quadratic polynomial where* $a_{i,j}$ *are pairwise different real parameters except that* $a_{i,j} = a_{j,i} (1 \leq i,j \leq n)$*. Let* $G(\boldsymbol{x}_n) = g(x_1^2, \ldots, x_n^2)$*, then* $\mathtt{Np}(G, [\boldsymbol{x}_n]) = \det(A_n)$.

Proof. By Theorem 50, we have

$$\mathtt{Np}(G, [\boldsymbol{x}_n], n) = \mathtt{Bp}(\det(A_{n-1})\det(A_n)x_n, x_n) = \det(A_{n-1})\det(A_n).$$

Therefore, $\mathtt{Np}(G, [\boldsymbol{x}_n]) = \gcd(\mathtt{Np}(G, [\boldsymbol{x}_n], 1), \ldots, \mathtt{Np}(G, [\boldsymbol{x}_n], n)) = \det(A_n)$. $\square$

By similar method we can prove that

Theorem 53. *Suppose* $g(\boldsymbol{x}_n) = \sum_{1\leq i,j\leq n} a_{i,j}x_ix_j = \boldsymbol{x}_n A_n \boldsymbol{x}_n^T$ *is a quadratic form where* $a_{i,j}$ *are pairwise different real parameters except that* $a_{i,j} = a_{j,i} (1 \leq i,j \leq n)$*. Let* $A_n = (a_{i,j})_{i,j=1}^n$*. Then* $\mathtt{Hp}(g, [\boldsymbol{x}_n]) = \det(A_n) = \operatorname{discrim}(g, [\boldsymbol{x}_n])$*, where* $\operatorname{discrim}(g, [\boldsymbol{x}_n])$ *is the discriminant of the quadratic form* g*, it is an irreducible polynomial in* $\mathbb{Z}[\boldsymbol{a}_{i,j}]$.

This theorem implies that, for a class of polynomial g, $\mathtt{Hp}(g, [\boldsymbol{x}_n])$ may coincide with its discriminant.

Theorem 49 and Theorem 50 show that, for a generic copositive problem, we can compute the projection set $\overline{\mathtt{Np}}(F, n-1) = \{f, \mathtt{Np}(f, [x_n]), \ldots, \mathtt{Np}(f, [x_n, \ldots, x_2])\}$ directly. Based on the theorem, it is easy to design a complete algorithm for solving copositive problems. However, for an input $f(\boldsymbol{x}_n)$, checking whether $f(\boldsymbol{x}_n)$ satisfies the hypothesis (3) of Theorem 49 is expensive. Therefore we propose a special incomplete algorithm `CMT` for **c**opositive **m**atrix **t**esting, which is formally described as Algorithm 7.

Remark 54. In Algorithm 7, we do not check the hypotheses of Theorem 49. Thus the algorithm is incomplete. However, the algorithm still makes sense because almost all $f(\boldsymbol{x}_n)$ defined by Eq. (3) satisfy the hypotheses. On the other hand, for an input $f(\boldsymbol{x}_n)$, checking whether $f(\boldsymbol{x}_n)$ satisfies the hypothesis (3) of Theorem 49 is expensive but the other three hypotheses are easy to check. Furthermore, $f(\boldsymbol{x}_n)$ is degenerate when some hypotheses do not hold and such case can be easily handled. Therefore, when implementing Algorithm `CMT`, we take into account those possible improvements. The details are omitted here.

Complexity analysis of Algorithm 7. We analyze the upper bound on the number of algebraic operations of Algorithm 7.

We first estimate the complexity of computing $\det(P_k)$ for $1 \leq k \leq n$. Because the entries of the last row and the last column of P_k are polynomials with k^2 terms and the other entries are integers, we expand $\det(P_k)$ by minors along the last column and then expand the minors again along the last rows. Therefore, the complexity of computing $\det(P_k)$, *i.e.*, g_k, is $\mathcal{O}(k^2(k-2)^3 + k^2(n-k)^2)$. Since $g_k(x_k, \boldsymbol{\alpha})$ is an even quartic univariate polynomial, the complexity of real root isolation for $g_k(x_k, \boldsymbol{\alpha})$ is $\mathcal{O}(1)$ and we only need to choose positive sample points when calling `SPOne`. That means $\mathtt{SPOne}(g_k(x_k, \boldsymbol{\alpha}), 1)$ returns at most 3 points. Thus the scale of O in line 13 is at most 3^{i-1}. The cost of

Algorithm 7. CMT

```
Input: An even quartic squarefree polynomial $F(\boldsymbol{x}_n) \in \mathbb{Z}[\boldsymbol{x}_n]$, $n \geq 1$, with an ordering
    $x_n \prec x_{n-1} \cdots \prec x_1$ and a set $Q$ of nonnegative polynomials.
Output: Whether or not $F(\boldsymbol{x}_n) \geq 0$ on $\mathbb{R}^n$
 1: if $F \in Q$ then return true
 2: end if
 3: for $i$ from 1 to $n$ do
 4:     if CMT($F(x_1, x_2, \ldots, x_n)|_{x_i=0}, Q$) =false then return false
 5:     else $Q := Q \cup (F(x_1, x_2, \ldots, x_n)|_{x_i=0})$
 6:     end if
 7: end for
 8: $g_n(x_n) := \det(P_n)$        ▷ Recall that $F(\boldsymbol{x}_n) = (x_1^2, \ldots, x_{n-1}^2, 1)P_n(x_1^2, \ldots, x_{n-1}^2, 1)^T$.
 9: $O :=$ SPOne($g_n, 1$)
10: for $i$ from 2 to $n$ do
11:     $S := \emptyset$
12:     $g_{n-i+1}(x_{n-i+1}, \ldots, x_n) := \det(P_{n-i+1})$
13:     for $\boldsymbol{\alpha}$ in $O$ do
14:         $S := S \bigcup (\boldsymbol{\alpha} \boxplus$ SPOne($g_{n-i+1}(x_{n-i+1}, \boldsymbol{\alpha}), 1$))
15:     end for
16:     $O := S$
17: end for
18: if $\exists \boldsymbol{\alpha}_n \in O$ such that $F(\boldsymbol{\alpha}_n) < 0$ then return false
19: end if
20: return true
```

computing $g_k(x_k, \boldsymbol{\alpha})$ is $\mathcal{O}(k^2)$ for each sample point $\boldsymbol{\alpha}$. Then the cost of the "for loop" at lines 10-17 is bounded by

$$\mathcal{O}(\sum_{i=1}^{n}(i^2(i-2)^3 + i^2(n-i)^2 + i^2 3^{i-1})) = \mathcal{O}(n^2 3^n).$$

In line 18 of Algorithm 7, the number of checking $F(\boldsymbol{\alpha}_n)$ is at most 3^n. And the complexity of every check in line 18 is $\mathcal{O}(n^2)$ since F has at most n^2 terms. Then, the complexity of line 18 is bounded by $\mathcal{O}(n^2 3^n)$. Therefore, the complexity of lines $8-19$ is bounded by $\mathcal{O}(n^2 3^n)$.

The scale of the set Q is at most $\sum_{k=0}^{n} \binom{n}{k} = 2^n$. So, the cost of all recursive calls is bounded by

$$\mathcal{O}(\sum_{k=0}^{n} \binom{n}{k} 3^k k^2) = \mathcal{O}(n^2 2^{2n}).$$

In conclusion, the complexity of Algorithm 7 is bounded by $\mathcal{O}(n^2 2^{2n})$.

Remark 55. By a more careful discussion, we may choose at most two sample points on every call `SPOne`($g_k(x_k, \boldsymbol{\alpha}), 1$). That will lead to an upper bound complexity, $\mathcal{O}(3^n n^2)$.

8. Examples

The Algorithm `HpTwo`, Algorithm `PSD-HpTwo`, and Algorithm `CMT` have been implemented as three programs using Maple. In this section, we report the performance of the three programs, respectively. All the timings in the tables are in seconds.

Example 9. (Strzeboński, 2000)

$$f = ax^3 + (a + b + c)x^2 + (a^2 + b^2 + c^2)x + a^3 + b^3 + c^3 - 1.$$

Under the order $a \prec b \prec c \prec x$, an open CAD defined by f has 132 sample points, while an open sample obtained by the algorithm `HpTwo` has 15 sample points.

Example 10. (Han et al., 2014)

$$f = x^4 - 2x^2y^2 + 2x^2z^2 + y^4 - 2y^2z^2 + z^4 + 2x^2 + 2y^2 - 4z^2 - 4.$$

Under the order $z \succ y \succ x$, an open CAD defined by f has 113 sample points, while an open sample obtained by the algorithm `HpTwo` has 87 sample points.

Example 11. For 100 random polynomials $f(x, y, z)$ with degree 8[2], Figure 3 shows the numbers of real roots of $\mathtt{Bp}(f, [z, y])$, $\mathtt{Bp}(f, [y, z])$ and $\mathtt{Hp}(f, [y, z])$, respectively. It is clear that the number of real roots of $\mathtt{Hp}(f, [y, z])$ is always less than those of $\mathtt{Bp}(f, [z, y])$ and $\mathtt{Bp}(f, [y, z])$.



Fig. 3. The number of real roots.

Example 12. In this example, we compare the performance of Algorithm `HpTwo` with open CAD on randomly generated polynomials. All the data in this example were obtained on a PC with Intel(R) Core(TM) i5 3.20GHz CPU, 8GB RAM, Windows 7 and Maple 17.

In the following table, we list the average time of projection phase and lifting phase, and the average number of sample points on 30 random polynomials with 4 variables and degree 4 generated by `randpoly([x,y,z,w],degree=4)-1`.

---

[2] Generated by `randpoly([x,y,z],degree=8)` in Maple 15.

	Projection	Lifting	Sample points
`HpTwo`	0.13	0.29	262
`open CAD`	0.19	3.11	486

If we get random polynomials with 5 variables and degree 3 by the command

$$\texttt{randpoly}([\texttt{seq}(\texttt{x}[\texttt{i}], \texttt{i} = 1..5)], \texttt{degree} = 3),$$

then the degrees of some variables are usually one. That makes the computation very easy for both `HpTwo` and open CAD. Therefore, we run the command $\texttt{randpoly}([\texttt{seq}(\texttt{x}[\texttt{i}], \texttt{i} = 1..5)], \texttt{degree} = 3) + \texttt{add}(\texttt{x}[\texttt{i}]^2, \texttt{i} = 1..5) - 1$ ten times to generate 10 random polynomials with 5 variables and degree 3. The data on the 10 polynomials are listed in the following table.

	Projection	Lifting	Sample points
`HpTwo`	2.87	3.51	2894
`open CAD`	0.76	12.01	7802

For many random polynomials with 4 variables and degree greater than 4 (or 5 variables and degree greater than 3), neither `HpTwo` nor open CAD can finish computation in reasonable time.

A main application of the new projection operator `Hp` is testing semi-definiteness of polynomials. Now, we illustrate the performance of our implementation of Algorithm `PSD-HpTwo` and Algorithm `CMT` with several non-trivial examples. For more examples, please visit the homepage[3] of the second author.

We report the timings of the programs `CMT`, `PSD-HpTwo`, and `DPS` (Han et al., 2013), the function PartialCylindricalAlgebraicDecomposition (`PCAD`) in Maple 15, function FindInstance (`FI`) in Mathematica 9, QEPCAD B (`QEPCAD`), the program `RAGlib`[4], and `SOSTOOLS` in MATLAB [5] on these examples.

`QEPCAD` and `SOSTOOLS` were performed on a PC with Intel(R) Core(TM) i5 3.20GHz CPU, 4GB RAM and ubuntu. The other computations were performed on a laptop with Inter Core(TM) i5-3317U 1.70GHz CPU, 4GB RAM, Windows 8 and Maple 15.

Example 13. (Han, 2011) Prove that

$$F(\boldsymbol{x}_n) = (\sum_{i=1}^{n} x_i^2)^2 - 4\sum_{i=1}^{n} x_i^2 x_{i+1}^2 \geq 0,$$

where $x_{n+1} = x_1$.

Hereafter "∞" means either the running time is over 4000 seconds or the software fails to get an answer.

---

[3] `https://sites.google.com/site/jingjunhan/home/software`

[4] `RAGlib` release 3.23 (Mar., 2015). The `RAGlib` has gone through significant improvements. Thus, we updated the timing using the most recent version.

[5] The MATLAB version is R2011b, SOSTOOLS's version is 3.00 and SeDuMi's version is 1.3.

n	5	8	11	17	23
`CMT`	0.06	0.48	1.28	4.87	11.95
`PSD − HpTwo`	0.28	0.95	6.26	29.53	140.01
`RAGlib`	0.42	0.76	1.34	3.95	8.25
`DPS`	0.29	∞	∞	∞	∞
`FI`	0.10	∞	∞	∞	∞
`PCAD`	0.26	∞	∞	∞	∞
`QEPCAD`	0.10	∞	∞	∞	∞
`SOSTOOLS`	0.23	1.38	3.94	247.56	∞

We then test the semi-definiteness of the polynomials (in fact, all $G(\boldsymbol{x}_n)$ are indefinite)

$$G(\boldsymbol{x}_n) = F(\boldsymbol{x}_n) - \frac{1}{10^{10}} x_n^4.$$

The timings are reported in the following table.

n	`CMT`	`PSD-HpTwo`	`RAGlib`	`DPS`	`FI`	`PCAD`	`QEPCAD`
20	1.81	3.828	0.59	∞	∞	∞	∞
30	5.59	13.594	2.01	∞	∞	∞	∞

Example 14. Prove that

$$B(\boldsymbol{x}_{3m+2}) = (\sum_{i=1}^{3m+2} x_i^2)^2 - 2\sum_{i=1}^{3m+2} x_i^2 \sum_{j=1}^{m} x_{i+3j+1}^2 \geq 0,$$

where $x_{3m+2+r} = x_r$. If $m = 1$, it is equivalent to the case $n = 5$ of Example 13. This form was once studied in Parrilo (2000).

m	`CMT`	`PSD-HpTwo`	`RAGlib`	`DPS`	`FI`	`PCAD`	`QEPCAD`
1	0.03	0.296	0.42	0.297	0.1	0.26	0.104
2	0.56	1.390	0.36	23.094	∞	∞	∞
3	0.71	9.672	0.75	∞	∞	∞	∞
4	7.68	∞	0.87	∞	∞	∞	∞

Remark 56. As showed by Example 12, according to our experiments, the application of `HpTwo` and `PSD-HpTwo` is limited at 3-4 variables and low degrees generally. It is not difficult to see that, if the input polynomial $f(\boldsymbol{x}_n)$ is symmetric, the new projection operator `Hp` cannot reduce the projection scale and the number of sample points. Thus, it is reasonable to conclude that the complexity of `PSD-HpTwo` is still doubly exponential.

Acknowledgements

Thank M. Safey El Din who provided us several examples and communicated with us on the usage of `RAGlib`.

The authors would like to convey their gratitude to all the four referees of our ISSAC'2014 paper, who provided their valuable comments, advice and suggestion, which helped improve this paper greatly on not only the presentation but also the technical details.

References

Andersson, L.-E., Chang, G., Elfving, T., 1995. Criteria for copositive matrices using simplices and barycentric coordinates. Linear Algebra and Its Applications 220, 9–30.

Basu, S., Pollack, R., Roy, M.-F., 1998. A new algorithm to find a point in every cell defined by a family of polynomials. In: Quantifier elimination and cylindrical algebraic decomposition. Springer, pp. 341–350.

Brown, C. W., 2001. Improved projection for cylindrical algebraic decomposition. J. Symb. Comput. 32 (5), 447–465.

Collins, G. E., 1975. Quantifier elimination for real closed fields by cylindrical algebraic decompostion. In: Automata Theory and Formal Languages 2nd GI Conference Kaiserslautern, May 20–23, 1975. Springer, pp. 134–183.

Collins, G. E., 1998. Quantifier elimination by cylindrical algebraic decomposition–twenty years of progress. In: Quantifier elimination and cylindrical algebraic decomposition. Springer, pp. 8–23.

Collins, G. E., Hong, H., 1991. Partial cylindrical algebraic decomposition for quantifier elimination. J. Symb. Comput. 12 (3), 299–328.

Hadeler, K., 1983. On copositive matrices. Linear Algebra and its Applications 49, 79–89.

Han, J., 2011. An Introduction to the Proving of Elementary Inequalities. Vol. 123. Harbin Institute of Technology Press.

Han, J., Dai, L., Xia, B., 2014. Constructing fewer open cells by gcd computation in cad projection. In: Proc. ISSAC'2014. ACM, pp. 240–247.

Han, J., Jin, Z., Xia, B., 2013. Proving inequalities and solving global optimization problems via simplified cad projection. arXiv:1205.1224.

Hiriart-Urruty, J.-B., Seeger, A., 2010. A variational approach to copositive matrices. SIAM review 52 (4), 593–629.

Hong, H., 1990. An improvement of the projection operator in cylindrical algebraic decomposition. In: Proc. ISSAC'1990. ACM, pp. 261–264.

Hong, H., Safey El Din, M., Jul. 2012. Variant quantifier elimination. J. Symb. Comput. 47 (7), 883–901.
URL `http://dx.doi.org/10.1016/j.jsc.2011.05.014`

McCallum, S., 1988. An improved projection operation for cylindrical algebraic decomposition of three-dimensional space. J. Symb. Comput. 5 (1), 141–161.

McCallum, S., 1998. An improved projection operation for cylindrical algebraic decomposition. In: Quantifier Elimination and Cylindrical Algebraic Decomposition. Springer, pp. 242–268.

Murty, K. G., Kabadi, S. N., 1987. Some np-complete problems in quadratic and nonlinear programming. Mathematical programming 39 (2), 117–129.

Parrilo, P. A., 2000. Structured semidefinite programs and semialgebraic geometry methods in robustness and optimization. Ph.D. thesis, CIT.

Safey El Din, M., Schost, É., 2003. Polar varieties and computation of one point in each connected component of a smooth real algebraic set. In: Proc. ISSAC'2003. ACM, pp. 224–231.

Strzeboński, A., 2000. Solving systems of strict polynomial inequalities. J. Symb. Comput. 29 (3), 471–480.